

University of Groningen

## Role of MYC in pediatric and adult B-cell lymphoma patients

Aukema, Sietse Marten

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Aukema, S. M. (2017). *Role of MYC in pediatric and adult B-cell lymphoma patients*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# 2

## General introduction to cancer cytogenetics of mature B-cell lymphomas (follicular lymphoma, diffuse large B-cell lymphoma & Burkitt lymphoma)

---

The following chapter is a selection of the published book chapter and a full version including other mature B-cell lymphoma entities, Hodgkin lymphoma and T-cell lymphomas has been published in:

Siebert R, Aukema SM Mature B- and T-cell Neoplasms and Hodgkin Lymphoma. Kapitel 11: 252-331, DOI: 10.1002/9781118795569.ch11 in Sverre Heim, Felix Mitelman (Eds). *Cancer Cytogenetics: Chromosomal and Molecular Genetic Aberrations of Tumor Cells*. 4. Edition (2015). Wiley-Blackwell, London.

Reproduced with permission.

Mature lymphoid neoplasms are an extremely heterogeneous group of malignancies with regard to biological, clinical, and morphological features (Jaffe et al., 2001; Swerdlow et al., 2008). The clinical presentation mainly includes leukemic forms and tumors affecting predominantly lymphatic organs, such as the lymph nodes or spleen, or nonlymphatic organs, such as the skin or central nervous system (CNS). Both solid and circulating phases are not infrequently present together in lymphoid neoplasms rendering distinction between “leukemia” and “lymphoma” artificial in many cases. The natural course run by the different diseases ranges from the rather indolent, like in some types of chronic lymphocytic leukemia (CLL) or follicular lymphoma (FL), to the highly aggressive, as exemplified by Burkitt lymphoma (BL).

Historically, the main division of mature lymphoid neoplasms has been between Hodgkin disease, nowadays called Hodgkin lymphoma (HL), and non-Hodgkin lymphoma (NHL). More than 99% of HL and approximately 85% of mature lymphoid neoplasms generally derive from lymphoid cells of the B-lineage, the remainder being derived from the T/natural killer (NK)-cell lineage. Considering this biological and clinical heterogeneity as well as the relatively easy access to samples from lymphatic malignancies, it is not surprising that a huge number of chromosomal aberrations have been described in mature lymphoid neoplasms and also that many of them are nonrandom or even specific (Mitelman et al., 2014). Since Manolov and Manolova (1972) first described a 14q + marker chromosome in BL and Zech et al. (1976) identified the marker as stemming from a t(8;14) (q24;q32), the first characteristic translocation to be identified in a malignant lymphoma, the karyotypes of several thousand mature lymphoid neoplasms have been made publicly available, and many of the recurrent aberrations have been molecularly characterized. Novel cryptic chromosomal rearrangements are continuously being identified using molecular cytogenetic approaches such as fluorescence *in situ* hybridization (FISH); microarray-based technologies, including array comparative genomic hybridization (aCGH); and lately also highthroughput sequencing (for reviews, see Martin-Subero et al., 2003a, 2003b; Schwaenen et al., 2003; Dewald et al., 2004; Kearney and Horsley, 2005; Ventura et al., 2006; Nieländer et al., 2007; Wolff et al., 2007; Kluin and Schuuring, 2011; Campo, 2013). The new information obtained via a variety of recently introduced techniques is constantly refining our picture of the chromosome aberrations of malignant lymphomas and chronic lymphoproliferative disorders, deepening our understanding of how molecular and chromosomal rearrangements are crucial events

in tumorigenesis. Nevertheless, the new technologies in no way render superfluous the cytogenetic analysis of banded chromosomes in metaphase plates: also, high-throughput sequencing suffers from limitations in, among other things, the ability to detect chromosomal fusions with the breakpoints in repetitive-rich sequences (e.g., Robertsonian translocations or rearrangements involving immunoglobulin loci). The new technological approaches refine our diagnostic abilities rather than supplant the time-honored, older techniques.

### Classification of mature lymphoid neoplasms

The identification and characterization of recurrent chromosomal changes in the tumor cells has influenced and continues to have a major impact on the classification of mature lymphoid neoplasms (Harris et al., 1994; Jaffe et al., 2001; Swerdlow et al., 2008; Kluin and Schuuring, 2011). For example, mantle cell lymphoma (MCL) was only accepted as a separate entity once the translocation t(11;14) (q13;q32) leading to cyclin D1 activation through the juxtaposition of *CCND1* to *IGH* was identified as a genetic hallmark of this disease (Banks et al., 1992).

A problem when assessing the many cytogenetic studies providing the data on which this chapter is based is that most of them were performed before the publication and widespread acceptance of the “WHO classification of tumors of hematopoietic and lymphoid tissues” (Jaffe et al., 2001; Swerdlow et al., 2008). Thus, in many instances, the diagnosis then given is no longer considered valid, or it may not even exist anymore; in fact, it is for this very reason that the title of the present chapter has been changed from that of the earliest editions of *Cancer Cytogenetics*. Usually, the new classification reflects an increased understanding of disease biology; for example, what was formerly called t(11;14)-positive CLL or B-cell prolymphocytic leukemia (B-PLL) is today, based on modern immunophenotyping analyses, mostly diagnosed as (leukemic) MCL or “MCL-like” disease (van der Velden et al., 2014), and cases of true t(11;14)-positive CLL or B-PLL are hardly recognized any longer. In addition, the detection of novel cytogenetic correlations and the increased use of new gene expression and sequencing technologies make regular revisions of the WHO classification necessary. The present chapter widely relies on the 2001 and 2008 WHO classifications (Swerdlow et al., 2008). However, as we write, a new update of the WHO classification is already being prepared. We therefore try to comment also on such genetic findings that are likely to be featured there.

The WHO classification groups lymphoid neoplasms according to four main criteria, namely, their clinical, morphologic, immunophenotypic, and genetic features. The relative diagnostic importance of these parameters differs among diseases; there is no one “gold standard” (Jaffe et al., 2001; Swerdlow et al., 2008). For each neoplasm, a cell of origin (COO) is postulated. Often, this putative COO represents the stage of differentiation rather than the cell in which the actual initial transforming event occurred, since the latter is not known with certainty. Therefore, the classification is widely based on how we perceive normal B- and T-cell developments to proceed.

The cytogenetic findings will be presented under the main headings “Mature B-cell neoplasms,” “Mature T-cell and NK-cell neoplasms,” and “HL.” For the convenience of the reader, at the beginning of each section, a tabular comparison of the most important relevant classification systems is provided.


**B- and T-cell development and physiological B- and T-cell receptor rearrangements**

The development and maturation of B and T lymphocytes are tightly regulated processes that take place in the bone marrow, peripheral blood, and lymphoid organs. Every maturation stage is characterized by physiological genetic rearrangements and expression of certain surface molecules and cytokines (for reviews, see Küppers et al., 1999; Küppers and Dalla-Favera, 2001; Küppers, 2005; Seifert et al., 2013).

The cells of the immune system are able to recognize multiple antigens through a multitude of different antigen-specific receptors. This diversity is generated through programmed intrachromosomal rearrangements of the genes encoding immunoglobulins (IG) in B lymphocytes and Tcell receptors (TCR) in T lymphocytes. These genes contain all or part of gene segments for variability (V), diversity (D), joining (J), and constant (C) sequences. Rearrangements of the V, D, and J genes are responsible for the great variability and specificity of the immune system.

Immunoglobulin molecules, also called antigen receptors or antibodies, are composed of Ig heavy chains (IgH) and light chains, either kappa (Igκ) or lambda (Igλ). These proteins are encoded by the *IGH* locus at chromosome band 14q32, *IGK* at 2p12, and *IGL* at 22q11. The *IGH* gene is composed of V, D, J, and C gene segments (Figure 2.1), whereas *IG* light-chain genes lack D segments. In

the bone marrow, B-cell development is initiated by rearrangements of *IGH* and *IGK/L* loci in B-cell progenitors. The so-called V(D)J in *IGH* (Figure 2.1) and VJ in *IGK/L* recombinations create a functional Ig. This Ig, of IgM isotype but also IgD in mature B cells, is expressed on the surface of B cells and is essential for cell development and survival. Those cells that produce nonfunctional rearrangements die by apoptosis. B cells leave the bone marrow to become mature, naive (not exposed to antigen) cells that travel through the bloodstream to secondary lymphoid organs such as the lymph nodes, spleen, and mucosa-associated lymphoid tissue (MALT). There, the B cells that interact with antigens through membrane-bound Ig are clonally selected and undergo a process of proliferation and differentiation to create a germinal center. The genomic DNA is then modified by receptor editing, class switching, and somatic hypermutation.



**Figure 2.1 |** Schematic representation of the mechanisms leading to *IGH* translocations in B-cell malignancies (based on Willis and Dyer, 2000; Küppers and Dalla-Favera, 2001). (A) *IGH* locus germline configuration displaying variable (VH), joining (JH), diversity (DH), and constant (C) regions. Switch regions (black and white rectangles) lie upstream of each constant region with the exception of Cδ. Eμ and Eα enhancers are shown in red. (B) Schematic representation of the VDJ recombination, class switch recombination, and somatic hypermutation mechanisms. (C) Formation of *IGH* translocations. Depending on the breakpoint location at 14q32, the *IGH* locus may deregulate proto-oncogenes in both partner chromosomes through the translocation (elbowed arrows).

Receptor editing is a process by which an originally expressed antibody polypeptide chain, usually a light chain, is replaced by another one. The process is mediated by secondary rearrangements of the variable region segments. Some B cells change their Ig isotype from IgM and IgD to IgG, IgA, or IgE, a process

called class switching (Figure 2.1). This event is mediated through further recombinations of the *IGH* locus, by which DNA between switch regions immediately upstream of the constant regions is deleted. Somatic hypermutation is the process by which mutations are introduced into the variable sequences of *IG* genes to increase the affinity for the antigen. Eventually, some of these cells with mutant antibodies expressing increased affinity are released to the peripheral blood as antibody-secreting cells, plasma cells, or long-lived memory cells.

The T-cell receptor is a heterodimer composed of either  $\alpha$  and  $\beta$  or  $\gamma$  and  $\delta$  chains. The genes encoding these proteins are *TCRA/D* located at 14q11, *TCRB* at 7q34, and *TCRG* at 7p14. The process by which these genes are assembled to make a functional T-cell receptor is similar to that occurring at *IG* loci in B cells. However, in contrast to B lymphocytes, T lymphocytes proliferate and differentiate along several different developmental pathways generating distinct subpopulations of mature T cells.

### Chromosomal translocations involving *IG* and *TCR* loci

*IG* loci, including both the *IGH* locus in 14q32 and the *IGK* and *IGL* light-chain loci in 2p12 and 22q11, respectively, are recurrently involved in many chromosomal translocations in mature B-cell malignancies (Table 2.1). In contrast, the *TCR* loci are rather seldom involved in structural aberrations in mature T-cell neoplasms with the notable exception of T-cell prolymphocytic leukemia (T-PLL), in which *inv(14)(q11q32)* and its variant *t(14;14)(q11;q32)* are present in approximately 80%, with some of the remaining cases showing the rare but recurrent *t(X;14)(q28;q11)* (discussed in the following).

The frequency of *IG* translocations in different subtypes of mature B-cell malignancies is quite variable. For some diseases, like BL with its *t(8;14)(q24;q32)* and light-chain variants or MCL with its *t(11;14)(q13;q32)*, one or a group of distinct *IG* translocations are pathognomonic for the disease and, hence, detectable in more than 95% of all cases (Fu et al., 2005; Hummel et al., 2006; Jares et al., 2007; Salaverria and Siebert, 2011). In other instances, like diffuse large B-cell lymphoma (DLBCL) or multiple myeloma (MM), *IG* translocations are detectable in roughly half of the cases but recombining with a large number of partners (Siebert et al., 2001; Kuehl and Bergsagel, 2002; Lenz et al., 2007; Martin et al., 2010; Walker et al., 2013). At the other end of the spectrum, CLL is notable for its lack of *IG* translocations, and when breaks do occur at 14q32 in this disease, they do not always involve the *IGH* locus (Nowakowski et al., 2005; Wlodarska et al., 2007).

*IG* translocations in malignant B cells may be multiple and may involve both *IGH* and a light-chain locus. A common combination is that of a *t(14;18)(q32;q21)* in FL being followed by a *t(8;14)* or a corresponding *t(2;8)* or *t(8;22)* variant at transformation to DLBCL (Knezevich et al., 2005; Martin-Subero et al., 2005). Such cases, known as “double-hit” lymphomas if they involve *IG* translocations to *MYC* and either *BCL2* or *BCL6*, are usually clinically aggressive and respond poorly to chemotherapy (Aukema et al., 2011). The latter example also shows that hardly any of the *IG* translocations in mature B-cell neoplasms is completely specific for a certain lymphoma subtype. For example, *t(11;14)(q13;q32)* is recurrent in both MCL and MM and *t(14;18)(q32;q21)* is recurrent in both FL and DLBCL but also recurrently occurs, in addition to trisomy 12, in CLL. The *t(8;14)(q24;q32)* and its variants can occur, besides in BL, in almost any kind of mature B-cell neoplasm, including CLL, MCL, FL, and DLBCL (Karsan et al., 1993; Au et al., 1999, 2000, 2002, 2004; Sen et al., 2002; Majid et al., 2007; Aukema et al., 2011, 2014; Horn et al., 2011; Put et al., 2012; de Oliveira et al., 2014).

Chromosomal translocations affecting *IG* loci that look identical at the cytogenetic level are also often identical in terms of which genes they target. In contrast, sometimes different partner genes are involved in seemingly identical translocations, like *BCL2* and *MALT1*, which recombine with *IGH* in *t(14;18)(q32;q21)* in FL and MALT lymphoma, respectively (Tsujimoto et al., 1985; Sanchez-Izquierdo et al., 2003; Streubel et al., 2003).

As outlined earlier, all *IG* loci are subject to multiple physiological breaks as part of recombination events during B-cell development. Chromosomal translocations are supposed to derive from mistakes in these recombination processes. Depending on whether mistakes occur in the V(D)J joining process, during class switching, or as by-products of somatic hypermutation, the chromosomal break-points are found in different regions of the *IG* loci (Figure 2.1). With regard to the heavy-chain locus, class switch-associated translocations affect the centromeric part of *IGH* in 14q32 containing the segments for the constant Ig regions, VDJs-associated translocations mostly affect the middle part of *IGH* that contains the J-segments, and somatic hypermutation-associated translocations predominantly affect the telomeric part of the *IGH* locus containing the V-segments (Willis and Dyer, 2000; Küppers and Dalla-Favera, 2001; Kuehl and Bergsagel, 2002; Dyer, 2003; Küppers, 2005).



The location of the IG breakpoint is supposed to reflect the developmental stage at which the translocation occurs in a B cell. In line with this hypothesis, the t(11;14)(q13;q32) of MCL, which is supposed to occur in a pregerminal center cell in the bone marrow through an unsuccessful *VDJ* rearrangement, mostly involves a J-segment. In contrast, the cytogenetically identical translocation in a subset of MM stems from a failed class switch recombination event supposed to take place in or after the germinal center and affects the region containing the constant genes (Walker et al., 2013). Similarly, IG translocations in FL are, with very rare exceptions (Fenton et al., 2002), supposed to derive from mistakes in the *VDJ*-recombination process, whereas those occurring in DLBCL, MM, or CLL mostly involve class switch regions (Willis and Dyer, 2000; Küppers and Dalla-Favera, 2001; Kuehl and Bergsagel, 2002; Dyer, 2003; Küppers, 2005). Remarkably, the t(8;14)(q24;q32) in endemic BL mostly involves the J, whereas the same translocation in sporadic BL mostly affects a constant region, indicating that different pathogenetic mechanisms are at work in the two situations (reviewed in Boxer and Dang, 2001; Küppers and Dalla-Favera, 2001).

The consequences of the IG translocations are activation of intact oncogenes through deregulation mediated by enhancer segments within the IG loci. Only on rare occasions are fusion transcripts of the IG and oncogenes generated (Willis and Dyer, 2000). Although *VDJ*-associated translocations of the *IGH* locus lead to activation of the oncogene on the der(14) through juxtaposition of the translocated oncogene with the *IGH* enhancer, class switch translocations in principle can activate oncogenes on both derivative chromosomes because the centromeric *IGH* enhancer(s) remains on der(14), whereas the telomeric *IGH* enhancer(s) translocates to the derivative partner chromosome (Figure 2.1). This explains why recurrent loss of the der(11) t(11;14) in MCL or der(14)t(4;14) in MM does not affect deregulation of the *CCND1* and *MMSET* genes in 11q13 and 4p16, respectively (Santra et al., 2003). In translocations affecting the *IGK* and *IGL* light-chain loci, the oncogenic activation usually takes place on the derivative partner chromosomes to which the enhancer elements were translocated (Willis and Dyer, 2000; Küppers and Dalla-Favera, 2001; Kuehl and Bergsagel, 2002; Dyer, 2003; Küppers, 2005; Kroenlein et al., 2012; Shimanuki et al., 2013).

**Table 2.1** | Chromosomal translocations and other structural aberrations affecting the *IGH* locus in 14q32, the *IGK* locus in 2p12, or the *IGL* locus in 22q11 in mature B-cell malignancies other than multiple myeloma

Translocation	Lymphoma/leukemia sub-type	Partner	References
t(1;14)(p22;q32)*	Mucosa-associated lymphoid tissue (MALT) lymphoma	<i>BCL10</i>	Willis et al. (1999) and Zhang et al. (1999)
t(1;14)(q21;32)	Diffuse large B-cell lymphoma	<i>MUC1</i>	Dyomin et al. (2000) and Gilles et al. (2000)
t(1;14)(q21;32)	Diffuse large B-cell lymphoma	<i>IRTA1</i>	Sonoki et al. (2004)
t(1;14)(q21;32)	Follicular lymphoma	<i>FCGR2B</i>	Chen et al. (2001a)
t(1;22)(q21;q11)	Follicular lymphoma	<i>FCGR2B</i>	Callanan et al. (2000)
t(2;7)(p11;q21)	Indolent B-cell lymphoproliferative disorders, splenic marginal zone lymphoma	<i>CDK6</i>	Corcoran et al. (1999) and Parker et al. (2011, 2013)
t(2;14)(p13;q32)	B-CLL, immunocytoma	<i>BCL11A</i>	Satterwhite et al. (2001)
t(3;14)(p14;q32)	MALT lymphoma, diffuse large B-cell lymphoma	<i>FOXP1</i>	Streubel et al. (2005)
t(3;14)(q27;q32)*	Diffuse large B-cell lymphoma, follicular lymphoma	<i>BCL6</i>	Kerckaert et al. (1993) and Ye et al. (1993a)
t(4;14)(q24;q32)	Posttransplant lymphoproliferative disorder	<i>BANK1</i>	Yan et al. (2014)
t(6;14)(p21;q32)*	Diffuse large B-cell lymphoma, other B-cell lymphomas	<i>CCND3</i>	Sonoki et al. (2001)
t(6;14)(p25;q32)	(Pediatric) diffuse large B-cell lymphoma/FL3B	<i>IRF4</i>	Salaverria et al. (2011)
t(6;14)(q15;q32)	Aggressive B-cell lymphoma, NOS	<i>BACH2</i>	Kobayashi et al. (2011)
t(7;14)(q21;q32)*	Splenic lymphoma with villous lymphocytes	<i>CDK6</i>	Corcoran et al. (1999)
t(8;14)(q24;q32)*	Burkitt lymphoma/leukemia, diffuse large B-cell lymphoma, B-cell prolymphocytic leukemia, multiple myeloma	<i>MYC</i>	Taub et al. (1982)
t(9;14)(p13;q32)	Lymphoplasmacytoid immunocytoma, other B-NHL	<i>PAX5</i>	Busslinger et al. (1996) and Iida et al. (1996)
t(9;14)(q33;q32)	Pediatric chronic myeloid leukemia (blast crisis)	<i>LHX2</i>	Nadal et al. (2012)
t(10;14)(q24;q32)	Diffuse large B-cell lymphoma	<i>NFKB2 (LYT10)</i>	Neri et al. (1991)
t(11;14)(p13;q32)	Diffuse large B-cell lymphoma, follicular lymphoma	<i>CD44</i>	Hu et al. (2012)

Translocation	Lymphoma/leukemia sub-type	Partner	References
t(11;14)(q13;q32)*	Mantle cell lymphoma, B-cell chronic lymphocytic leukemia, B-cell prolymphocytic leukemia, splenic lymphoma with villous lymphocytes	CCND1 (BCL1)	Motokura et al. (1991) and Rosenberg et al. (1991)
t(11;14)(q23;q32)	Primary mediastinal B-cell lymphoma	PAFAH1A2	Lecointe et al. (1999)
t(11;14)(q23;q32)	Diffuse large B-cell lymphoma	RCK	Akao et al. (1992)
t(12;14)(p13;q32)*	Mantle cell lymphoma	CCND2	Gesk et al. (2006) and Herens et al. (2008)
t(12;14)(q24;q32)	Burkitt lymphoma/leukemia, multiple myeloma	BCL7A	Zani et al. (1996)
del(14)(q24q32)	Chronic lymphocytic leukemia, low-grade B-NHL	ZFP36L1	Pospisilova et al. (2007) and Nagel et al. (2009a)
t(14;15)(q32;q11-13)	Diffuse large B-cell lymphoma	BCL8	Dyomin et al. (1997)
t(14;16)(q32;q24)	B-ALL (L3), aggressive B-cell lymphoma, NOS	CBFA2T3/ACSF3	Salaverria et al. (2012)
t(14;16)(q32.33;q24.1)	Diffuse large B-cell lymphoma	IRF8	Bouamar et al. (2013) and Tinguely et al. (2014)
t(14;18)(q32;q21)*	Follicular lymphoma, diffuse large B-cell lymphoma	BCL2	Tsujimoto et al. (1985)
t(14;18)(q32;q21)	Marginal zone lymphoma	MALT1	Sanchez-Izquierdo et al. (2003) and Streubel et al. (2003)
t(14;19)(q32;q12)	Diffuse large B-cell lymphoma	CCNE1	Nagel et al. (2009b)
t(14;19)(q32;q13)	Diffuse large B-cell lymphoma	SPIB	Lenz et al.-2007
t(14;19)(q32;q13)*	B-cell chronic lymphocytic leukemia, other B-NHL	BCL3	Ohno et al. (1990)
t(14;22)(q32;q11)	Low-grade B-NHL	IGL?	Aamot et al. (2005)
t(X;14)(p11;q32)	MALT lymphoma	GPR34	Ansell et al. (2012)

Modified from Willis and Dyer (2000), Siebert et al. (2001), and Dyer (2013). Only those translocations are shown in which the partner genes have been identified by molecular cloning or equivalent technique(s). Several other IG translocations have been identified and in part FISH mapped, like t(14;19)(q32;p13) (Micci et al., 2007), but await unambiguous identification of the partners. \*Variants with IG light-chain loci in 2p12 and 22q11 have been described.

The breakpoint on the derivative partner chromosome can be located 3' or 5' of the target oncogene because the transforming mechanism is deregulated ex-

pression through aberrant juxtaposition with the IG enhancer rather than creation of a gene fusion or promoter substitution. Enhancers can exert their activity over distances of several hundred kilobases. In BL, for example, the breakpoints affecting the *MYC* gene can be located up to almost 1 Mb centromeric of *MYC* in t(8;14) and telomeric of *MYC* in t(8;22) and t(2;8) (Joos et al., 1992a, 1992b; Einerson et al., 2006; Kroenlein et al., 2012). Remarkably, the breakpoints of the t(8;14) in 8q24 vary between endemic and sporadic BL with the former more frequently affecting the far 5' (centromeric) part of the *MYC* locus (reviewed in Boxer and Dang, 2001; Küppers and Dalla-Favera, 2001). This variability of the breakpoints, which can also be observed in *MYC* translocations affecting non-IG loci, can pose considerable diagnostic problems if FISH is applied for their detection (Bertrand et al., 2007; May et al., 2010).

Gene deregulation may occasionally be a consequence of an insertion rather than a translocation. Thus, both *BCL2* and *CCND1* have been observed to be inserted into the *IGH* locus in cases of FL and MCL in rearrangements that functionally mimic the t(14;18) and t(11;14), respectively (Vaandrager et al., 2000; Fenton et al., 2004). Although the biological consequences of such insertions appear to be identical to those of regular translocations, their pathogenetic origins would appear to involve different DNA breaks.

Exceptions to the general rule that the recurrent translocations of mature lymphoid neoplasms rarely give rise to fusion genes occur predominantly in MALT lymphoma and T-cell lymphomas. In the former, the specific t(11;18)(q21;q21) leads to *API2-MALT1* fusion (Akagi et al., 1999; Dierlamm et al., 1999). In the latter, t(2;5)(p23;q35) as well as its rare variants lead to fusion of the anaplastic lymphoma kinase (*ALK*) gene in 2p23 with the nucleophosmin (*NPM*) gene or, in the case of variants, other partners (Morris et al., 1994). Finally, rare cases of peripheral T-cell lymphoma with an *ITK-SYK* fusion due to a t(5;9)(q33;q22) have been reported (Streubel et al., 2006b). Recent data from high-throughput sequencing studies suggest that a number of rare fusion genes remain to be discovered in lymphatic neoplasms.

The most common rearrangement of a non-IG locus in mature lymphoid neoplasms is the targeting of the *BCL6* gene in 3q27 that takes place in a variety of translocations with a multitude of chromosomal partners (Ueda et al., 2002a; Ohno, 2006) (Table 2.2). Common to all these translocations is that they lead to promoter substitution and, consequently, deregulation of the *BCL6* gene (Ohno,

2004, 2006; Jardin et al., 2007). Other oncogenes known to be targets of IG translocations are sometimes juxtaposed directly next to each other through translocations, such as *PAX5* and *MYC* through a t(8;9)(q24;p13) and *BCL6* and *MYC* through t(3;8)(q27;q24) in t(14;18)-positive FL (Bertrand et al., 2007; Sonoki et al., 2007; Johnson et al., 2009; Aukema et al., 2011). How the supposed *MYC* deregulation in these translocations occurs is so far unknown though an enhancer mechanism seems likely. There is no evidence for interposition of IG sequences like in t(8;14;18) (q24;q32;q21), a recurrent aberration in FL progression, in which both *MYC* and *BCL2* are deregulated from a common IG fragment between them (Dyer et al., 1996; Knezevich et al., 2005).

**Table 2.2 |** Chromosomal translocations affecting the *BCL6* locus in 3q27 in mature B-cell neoplasms

Translocation	Partner locus	<i>BCL6</i> partner gene	References
t(1;3)(q25;q27)	1q25	<i>GAS5</i>	Nakamura et al. (2008)
t(2;3)(p12;q27)	2p12	<i>IGK</i>	Bastard et al. (1992), De-weindt et al. (1993), and Akasaka et al. (2000a, 2000b)
t(3;3)(q25;q27)	3q25	<i>MBNL1</i>	Akasaka et al. (2003)
	3q26	<i>SIAT1</i>	Akasaka et al. (2003)
t(3;3)(q27;q27)	3q27	<i>ST6GAL1</i>	Akasaka et al. (2003)
t(3;3)(q27;q27)	3q27	<i>EIF4A2</i>	Yoshida et al. (1999) and Akasaka et al. (2003)
t(3;3)(q27;q29)	3q29	<i>TFRC</i>	Yoshida et al. (1999), Akasaka et al. (2000a, 2000b), and Chen et al. (2006)
t(3;4)(q27;p11–13)	4p11–13	<i>RhoH/TTF</i>	Dallery et al. (1995) and Akasaka et al (2000a, 2000b), (2003)
t(3;6)(q27;p22)	6p21–22	<i>HIST1H4I</i>	Akasaka et al. (1997), (2000a, 2000b), Kurata et al. (2002), and Schwindt et al. (2006)
t(3;6)(q27;p21)	6p21	<i>PIM1</i>	Yoshida et al. (1999), Akasaka et al.(2000a, 2000b), and Kaneita et al. (2001)
t(3;6)(q27;p21)	6p21	<i>SFRS3/SRP20</i>	Chen et al. (2001b) and Ohno (2006)
t(3;6)(q27;q15)	6q15	<i>U50HG/SNHG5</i>	Tanaka et al. (2000) and Akasaka et al. (2003)

t(3;7)(q27;p12)	7p12	<i>IKZF1</i>	Hosokawa et al. (2000) and Kaneita et al. (2001)
t(3;7)(q27;q32)	7q32	<i>MIR29/FRA7H</i>	Schneider et al. (2008)
t(3;8)(q27;q24.1)	8q24	<i>MYC</i>	Bertrand et al. (2007) and Sonoki et al. (2007)
t(3;9)(q27;p24)	9p24	<i>DMRT1</i>	Chen et al. (2006)
t(3;9)(q27;p11)	9p11	<i>GRHPR</i>	Akasaka et al. (2003)
t(3;11)(q27;q23)	11q23	<i>BOB1/OBF1/POU2AF1</i>	Galiègue-Zouitina et al. (1996) and Ohno (2006)
t(3;12)(q27;p13)	12p13	<i>GAPDH</i>	Montesinos-Rongen et al. (2003) and Chen et al. (2006)
t(3;12)(q27;q12)	12q12	<i>LRMP</i>	Akasaka et al. (2003)
t(3;12)(q27;q23–24)	12q23	<i>NACA</i>	Akasaka et al. (2000a, 2000b)
t(3;13)(q27;q14)	13q14	<i>LCP1</i>	Galiègue-Zouitina et al. (1999)
t(3;14)(q27;q32)	14q32	<i>IGH</i>	Bastard et al. (1992), De-weindt et al. (1993), Kerckaert et al. (1993), Ye et al. (1993b), and Akasaka et al. (2003)
t(3;14)(q27;q32)	14q32	<i>HSP89A</i>	Akasaka et al. (2000a, 2000b), Xu et al. (2000), Montesinos-Rongen et al. (2003), and Chen et al. (2006)
t(3;16)(q27;p13)	16p13	<i>CIITA</i>	Yoshida et al. (1999, Kaneita et al. (2001), and Akasaka et al. (2003)
t(3;16)(q27;p11)	16p11	<i>IL21R</i>	Ueda et al. (2002b)
t(3;19)(q27;q13)	19q13	<i>NAPA</i>	Akasaka et al. (2003)
t(3;22)(q27;q11)	22q11	<i>IGL</i>	Bastard et al. (1992), De-weindt et al. (1993), Miki et al. (1994b), and Schimanuki et al. (2013)

Data from <http://atlasgeneticsoncology.org/Genes/BCL6ID20.html>, Ueda et al. (2002a), and Ohno (2006). Chromosomal bands of *BCL6* partner genes and gene annotations may in individual cases be mapped to different chromosomal bands in various publications

Cytogenetic evolution in lymphoid neoplasms

Several cytogenetic and subsequent molecular genetic studies have led to the concept that the neoplastic process in many mature lymphoid malignancies follows a multistep model similar to that proposed for colorectal cancer and other malignancies. It is widely assumed that a primary genetic alteration of a tumor clone



initiates lymphomagenesis. Primary alterations are usually present in the stemline of a tumor. Chromosomal translocations, like several of the *IG* and *TCR* translocations, are most common as primary genetic alterations and can serve as valuable diagnostic markers. However, it should be noted that what in one disease setting may constitute a primary change in another may occur as a secondary abnormality.

As shown in mouse transgenic models, the primary genetic alteration is often not sufficient to drive lymphomagenesis (reviewed in Willis and Dyer, 2000). Moreover, by sensitive molecular techniques, several primary genetic alterations like t(14;18), t(11;14), or t(2;5) can also be detected in healthy individuals, sometimes at surprisingly high levels (Limpens et al., 1995; Maes et al., 2001; Bäsecke et al., 2002; Biagi and Seymour, 2002; Roulland et al., 2004, 2014). Although our understanding is limited as to what extent the cells carrying the translocations are susceptible to neoplastic transformation, it has been shown that the presence of t(14;18)-positive precursors is predictive for the later development of FL and the precursors can actually progress into FL (Roulland et al., 2014). These findings have contributed to the view that additional genetic changes are required for the full neoplastic phenotype of a mature lymphoid malignancy to emerge. The aberrations that bring about this phenotypic transformation are usually called secondary genetic changes. Consequently, the vast majority of lymphomas with a defined primary change, like t(14;18)-positive FL or t(11;14)-positive MCL, carry also secondary alterations (Johansson et al., 1995). The so-called tertiary genetic events may occur in some lymphomas and are associated with the transformation of an indolent malignancy to more aggressive forms. As will be described subsequently, secondary and tertiary chromosomal alterations in lymphoid malignancies predominantly lead to gain or loss of genetic material, whereas recurrent, balanced translocations—with the exception of translocations affecting the *MYC* locus in chromosome band 8q24—seem to be uncommon as secondary events. The number of secondary genetic aberrations, be they driver or passenger mutations, usually increases during disease progression (Knight et al., 2012; Beà et al., 2013; Landau et al., 2013; Bauer et al., 2014). As a rule of thumb, within a given lymphoma subtype, the prognosis is inversely related to the complexity of the karyotype or the extent of the secondary changes.

Secondary genetic aberrations may be present in only a subset of cells of a given tumor, and different secondary aberrations may be present in different subclones that may become the dominant clone during disease progression or at relapse (Knight et al., 2012; Landau et al., 2013; de Rossi et al., 2014). Tumors of the

same type (and harboring the same primary event) might thus show variation in terms of secondary aberrations. Nevertheless, the pattern of secondary changes in mature lymphoid neoplasms is not random but rather seems to depend on the nature of the primary event. Some primary genetic alterations like t(11;18)(q21;q21) are hardly ever associated with cytogenetically detectable secondary aberrations (Horsman et al., 1992; Ott et al., 1997; Barth et al., 2001). Other translocations such as t(1;14)(p22;q32) are associated with a highly conserved set of secondary events, including trisomy of chromosomes 3, 12, and 18. FL with t(14;18)(q32;q21) show different, distinctive patterns of secondary changes suggesting different pathways of germinal center B-cell transformation (Höglund et al., 2004). T-PLL is characterized by an extraordinary conservation of both primary and secondary genetic changes (reviewed by Dürig et al., 2007).

The mutational pattern in a given lymphoma might not only depend on the primary aberration but also on the mutational mechanisms active in this particular case (Alexandrov et al., 2013). Several mutational mechanisms have been shown to be operative in lymphomas contributing to the mutational landscape of the tumors. Indeed, the genome of a lymphatic neoplasm usually contains between 1000 and several thousand mutations (Puente et al., 2011; Richter et al., 2012; Alexandrov et al., 2013). Based on the mutational patterns detected, it has been shown for FL that disease progression can occur from the initial tumor or from a clonal progenitor not found at initial diagnosis (Okosun et al., 2014; Pasqualucci et al., 2014). Remarkably, at least in FL, chromosomal, molecular, and epigenetic changes seem to evolve, concordantly suggesting that a chromatin-acting factor like *CREBBP* or *MLL2* may be one of the regulators of clonal evolution (Loeffler et al., 2015). Tumor heterogeneity and its impact on progression and clinical outcome will become a major topic for future studies in lymphoma genetics (Knight et al., 2012; Schuh et al., 2012; Landau et al., 2013; Rossi et al., 2014; Suguro et al., 2014).

### **Chromosomal aberrations common to multiple subtypes of mature lymphoid neoplasms.**

In the following sections, the most common cytogenetic changes in mature lymphoid neoplasms will be discussed in the context of the disease subtypes in which they are most prominent. It should nevertheless be emphasized that many aberrations are recurrent in several or even all kinds of mature lymphoid neoplasms. These might constitute primary aberrations in some of them but are more likely to be secondary or even tertiary aberrations in the majority. Probably, the best example of such a low-specific change is deletion of 6q.

*Deletions of the long arm of chromosome 6* are frequent in all types of mature lymphoid neoplasms including HL. The incidence in different studies varies from roughly 5% in CLL to 30% in DLBCL up to even higher frequencies in T/NK-cell disorders. The deletions have mostly been interpreted as terminal. Band 6q21 seems to be the one most often affected. Offit et al. (1993) reviewed the chromosome 6 abnormalities in 459 consecutively ascertained karyotypically abnormal cases of NHL and suggested that three minimally common deleted regions could be discerned: a distal one encompassing 6q25–27, a second one involving the more proximal band 6q21, and the third one located between the two others involving band 6q23. Several loss of heterozygosity (LOH), molecular cytogenetic, and aCGH studies have meanwhile narrowed down critical regions in 6q to 6q21, 6q23.3, and 6q27. These studies have identified recurrent inactivation of the *PRDM1/BLIMP1* gene in 6q21 in activated B-cell-type (ABC-DLBCL) and T-cell lymphomas (Pasqualucci et al., 2006; Tam et al., 2006; Iqbal et al., 2009). Moreover, a minimally common deleted region in 6q23.3 containing the *TNFAIP3 (A20)* and *PERP* genes has been identified in FL and ocular adnexal marginal zone B-cell lymphoma (MZL), and mutations inactivating *TNFAIP3* have been detected in classical Hodgkin lymphoma (cHL) and primary mediastinal B-cell lymphomas (PMBCL) (Ross et al., 2007; Giefing et al., 2008; Honma et al., 2008; Schmitz et al., 2009).

The descriptions in this chapter will focus on chromosomal changes as they are detected by chromosome banding techniques. It needs to be stressed that the “real” frequencies of chromosomal aberrations may be underestimated if this approach is the only one taken, without help from molecular cytogenetic techniques like FISH and CGH, which can also be used to examine paraffin-embedded samples. Therefore, though the contribution of these techniques is not the prime concern of this book, data thus obtained will often be referred to as they have contributed mightily to the current picture of the cytogenetic changes of mature lymphoid neoplasms. For reviews focusing on the application of FISH and CGH in lymphoid neoplasms, the reader may be well advised to consult Martin-Subero et al. (2003a), Dewald et al. (2004), Kearney and Horsley (2005), Ventura et al. (2006), Wolff et al. (2007), and Kluin and Schuurin (2011). Finally, the exponentially fast-growing number of studies using aCGH and single-nucleotide polymorphism (SNP) chip-based technologies as well as next-generation sequencing will not be discussed in detail here. A review of the findings obtained by array-based technologies was published by Nieländer et al. (2007), and databasing of the findings has been initiated (<http://www.progenetix.net/>). For mutational patterns derived

from highthroughput sequencing, the reader is referred to constantly updated databases such as COSMIC (<http://cancer.sanger.ac.uk/cancergenome/projects/cosmic/>) and the reports from the ICGC project (<http://www.icgc.org/>).

### Mature B-cell neoplasms

A comparative overview of the different subtypes of mature B-cell neoplasms recognized by the Kiel, REAL, and WHO classifications is provided in Table 2.3. The clinically most relevant chromosomal aberrations are summarized in Table 2.4.

### FL (including pediatric FL and primary cutaneous follicle center lymphoma)

FL comprises 22% of all lymphomas worldwide but shows significant geographic variation. FL is rare in children, and the median age of incidence is 59 years. It is a neoplasm resembling follicle center B cells, that is, centrocytes and centroblasts. There is heterogeneity with regard to growth pattern (follicular, follicular and diffuse, and minimally follicular) and the proportion of centroblasts, the latter being reflected in three different grades (FL grades 1–3) with grade 1 showing the lowest numbers of centroblasts. Grade 3 shows the highest number of centroblasts and is subdivided into grade 3a in which centrocytes are present and grade 3b with solid sheets of centroblasts (Jaffe et al., 2001; Swerdlow et al., 2008). Variants include diffuse follicle center lymphoma and cutaneous follicle center lymphoma. Particularly, if considering the prominence of centroblasts in grade 3b and variants with diffuse growth patterns, it is not surprising that at this end of the morphologic spectrum, FL shows considerable overlap with DLBCL (Jaffe et al., 2001; Swerdlow et al., 2008). Moreover, there is a steady rate (3% per year) of transformation of FL mostly to DLBCL (Montoto et al., 2007; Montoto and Fitzgibbon, 2011). Finally, FL and DLBCL might coexist in a patient even in the same lymph node, rendering a clear distinction between the karyotypic features of the two diseases difficult. Approximately 80–90% of FL carry a *t(14;18) (q32;q21)* (Figure 2.2) or very seldom one of its variants, *t(2;18) (p12;q21)* or *t(18;22)(q21;q11)* (Johansson et al., 1995; Hillion et al., 1991; Bentley et al., 2005; Lin et al., 2008). It is noteworthy that the *t(14;18)* is extremely rare in FL patients below 18 years of age and virtually absent in children below the age of 14 years (Oschlies et al., 2010; Louissaint et al., 2012; Liu et al., 2013).

**Table 2.3 | Comparison of the Kiel, REAL, and WHO classifications of B-cell malignancies\***

<i>Kiel classification</i>	<i>REAL classification</i>	<i>WHO (2001)</i>	<i>WHO (2008)</i>
Peripheral B-cell neoplasms	Mature (peripheral) B cell neoplasms		
B-lymphocytic, B-CLL, B-PLL	B-CLL/B-PLL/small lymphocytic lymphoma	<b>B-CLL</b> /small lymphocytic lymphoma	<b>B-CLL</b> /small lymphocytic lymphoma
Lymphoplasmacytoid immunocytoma	Variant B-CLL: with plasmacytoid differentiation	Variant B-CLL: with monoclonal gammopathy/ plasmacytoid differentiation	Variant B-CLL: with monoclonal gammopathy/ plasmacytoid differentiation
		B-PLL	B-PLL
Lymphoplasmacytic immunocytoma	Lymphoplasmacytoid lymphoma	Lymphoplasmacytoid lymphoma	Lymphoplasmacytic lymphoma
Centrocytic	Mantle cell lymphoma	Mantle cell lymphoma	Mantle cell lymphoma
Centroblastic,		Variant: blastoid	Morphologic variants:
centrocytoid subtype			blastoid, pleomorphic, small cell, marginal zone-like
Centroblastic–centrocytic, follicular	Follicular center lymphoma, follicular grades I and II	<b>Follicular lymphoma, grades I and II</b>	<b>Follicular lymphoma, grades I and II</b>
Centroblastic–centrocytic, diffuse	Follicular center lymphoma, diffuse, small cell*		
Centroblastic, follicular	Follicular center lymphoma, follicular grade III	Follicular lymphoma, grade III	Follicular lymphoma, grade III
		Cutaneous follicle center lymphoma	Cutaneous follicle center lymphoma
	Extranodal marginal zone B-cell lymphoma (low-grade B-cell lymphoma or MALT)	<b>Extranodal marginal zone B-cell lymphoma of MALT type</b>	<b>Extranodal marginal zone B-cell lymphoma of MALT type</b>
Monocytoid lymphoma, including marginal zone lymphoma	Nodal marginal zone B-cell lymphoma*	Nodal marginal zone B-cell lymphoma*	Nodal marginal zone B-cell lymphoma*

<b>Kiel classification</b>	<b>REAL classification</b>	<b>WHO (2001)</b>	<b>WHO (2008)</b>
	Splenic marginal zone B-cell lymphoma*	Splenic marginal zone B-cell lymphoma*	Splenic marginal zone B-cell lymphoma*
Hairy cell leukemia	Hairy cell leukemia	Hairy cell leukemia	Hairy cell leukemia
Plasmacytic	Plasmacytoma/myeloma	<b>Plasma cell myeloma/plasmacytoma</b>	<b>Plasma cell myeloma/plasmacytoma</b>
Centroblastic	Diffuse large B-cell lymphoma	<b>Diffuse large B-cell lymphoma</b> , variants: centroblastic, immunoblastic, T cell/histiocyte rich, lymphomatoid granulomatosis type, anaplastic large B-cell, plasmablastic	<b>Diffuse large B-cell lymphoma, NOS</b>
(monomorphic, polymorphic, and multilobated subtypes)			<p>Morphologic variants:</p> <p>centroblastic, immunoblastic, anaplastic</p> <p><b>Subtypes:</b> T cell/histiocyte rich</p> <p>Primary CNS</p> <p>Primary cutaneous, leg type</p> <p>EBV-positive DLBCL of the elderly</p> <p><b>Other:</b> DLBCL associated with chronic inflammation</p> <p>Lymphomatoid granulomatosis type</p> <p>ALK+ DLBCL</p> <p>HHV8-</p> <p>Associated multicentric Castleman disease</p>
B-immunoblastic			

Kiel classification	REAL classification	WHO (2001)	WHO (2008)
B-large cell anaplastic lymphoma	Primary mediastinal large B-cell lymphoma	Subtypes: mediastinal (thymic) large B-cell, primary effusion lymphoma, intravascular large B-cell lymphoma	Primary mediastinal (thymic) large B-cell lymphoma  Primary effusion lymphoma Intravascular large B-cell lymphoma
Burkitt lymphoma	Burkitt lymphoma  High-grade B-cell lymphoma, Burkitt-like*	Burkitt lymphoma  Burkitt cell leukemia  Atypical Burkitt lymphoma	Burkitt lymphoma  Burkitt cell leukemia/BALL L3  <b>B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and BL</b>

Data from Lennert et al. (1975), Harris et al. (1994), Jaffe et al. (2001), and Swerdlow et al. (2008).  
Bold letters represent the most common subtypes.  
\*Provisional entity.

Pediatric-type FL is recognized as a distinct but provisional entity in the 2008 WHO classification (Harris et al., 2008; Swerdlow et al., 2008) and is characterized by the absence of *t(14;18)(q32;q21)* and *BCL6* rearrangements and an overall indolent clinical behavior (Oschlies et al., 2010; Louissaint et al., 2012; Attarbaschi et al., 2013; Martin-Guerrero et al., 2013). A subset of cases (mainly FL3 with or without DLBCL component) shows *IG-IRF4* translocations with frequent chromosomal imbalances (Salaverria et al., 2011, 2013). In FL lacking *IG-IRF4* translocations, two subgroups may be distinguished: one “genetically normal” group with no DLBCL component and no copy number abnormalities and a “genetically aberrant group” (often with a DLBCL component) with more frequent copy number abnormalities, LOH at 1p36, and recurrent *TNFRSF14* and *EZH2* mutations (Salaverria et al., 2011; Martin-Guerrero et al., 2013). Pediatric-type FL is also encountered in young adults; then too it is characterized by a high prolifer-

ation rate but indolent clinical behavior (Louissaint et al., 2012; Liu et al., 2013).

In primary cutaneous follicle center lymphoma, conflicting data on the frequency of *t(14;18)* have been published. Variations among studies may be related to differences among patient cohorts and technique used (PCR vs. FISH) to detect the translocation; however, if any real frequency differences exist at all, the translocation seems to be less common in cutaneous than in nodal FL (Mirza et al., 2002; Vergier et al., 2004; Kim et al., 2005; Streubel et al., 2006a; Belaud-Rotureau et al., 2008; Gulia et al., 2011; Abdul-Wahab et al., 2014).

In nodal FL, there is evidence that the *t(14;18)* is more frequent in FL grades 1 and 2 (88% according to Katzenberger et al., 2004) than in FL 3, particularly FL 3b. Horn et al. (2011) identified using FISH *BCL2* breaks in 22/25 (88%), 7/12 (58%), and 2/23 (9%) of FL1/2, FL3A, and FL3B, respectively. Ott et al. (2002) found *t(14;18)* in 8 of 11 (73%) FL3a but only in 2 of 16 (13%) FL3b with or without a DLBCL component. Similarly, Katzenberger et al. (2004) found *t(14;18)* in only 1 of 27 FL3b (4%). Bosga-Bouwer et al. (2003) claimed the existence of three distinctive subgroups of FL3b based on the presence of breakpoints in 3q27, a translocation *t(14;18)*, or the absence of both. None of their FL grade 3B cases harbored both a *t(14;18)* and a 3q27 aberration. Indeed, *trisomy 3* and *breakpoints in 3q27* affecting the *BCL6* locus are the most prominent findings in *t(14;18)*-negative FL. As there is convincing evidence for a biological difference between *t(14;18)*-positive and *t(14;18)*-negative FL, these two groups will in the following be treated separately (Leich et al., 2009).

The nonrandom occurrence of *t(14;18)(q32;q21)* in NHL was first pointed out by Fukuhara et al. (1979). Yunis et al. (1984) mapped the breakpoints to subbands 14q32.3 and 18q21.1. The *t(14;18)* has since turned out to be the most common translocation in NHL (15% of the total) but with considerable geographic differences (Biagi and Seymour, 2002). When tested for molecularly, the translocation can also be detected in tissues from healthy donors to some extent depending on their age and external factors such as smoking habits and exposure to pesticides (Limpens et al., 1995; Biagi and Seymour, 2002; Roulland et al., 2004; Agopian et al., 2009; Tellier et al., 2014). *t(14;18)* is not restricted to FL but may also be detected in 30% of DLBCL. In that lymphoma, the translocation seems to be largely restricted to the germinal center B-cell-like type (GCB) as defined by gene expression profiling (Rosenwald et al., 2002; Visco et al., 2013). The frequent




**Table 2.4** | Overview of frequent and diagnostically relevant chromosomal aberrations in the most common mature B-lymphoid neoplasms

Neoplasm	Cytogenetic aberration	Frequency (%)
B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma	del(13q)	~50
	+12	~15
	del(11q)	15–20
Mantle cell lymphoma	del(6q), del(17p)	~5
	t/der(14)(q32)	4–20
	t(11;14)(q13;q32)	>95
	del(11q), del(13q), der(3q) +12, del(1p), del(6q), del(9p), del(17p)	10–50 5–15
Extranodal marginal zone B-cell lymphoma of MALT type	t(14;18)(q32;q21) involving MALT1	11
	t(3;14)(p14;q32)	<10
	t(1;14)(p22;q32)	<2
	+3/3q, +18/18q	30
Follicular lymphoma	t(14;18)(q32;q21) and variants involving BCL2	80–90
	t(3q27)	<10 (FL1/2)–55 (FL3b)
	+X, +7, +12/12q, +18/18q, +der(18)	
	t(14;18), del(6q), del(10q), del(17p), dup(1q), der(1p)	>10
Diffuse large B-cell lymphoma	t(3q27)	20–40
	t(14;18)(q32;q21)	20–30 (GCB)
	t(8;14)(q24;q32)/t(8q24)	5–15 (GCB > ABC)
	+3/3q, +18/18q, +19q, del(6q), del(9p)	10–40 (ABC)
	+1q, +2p13–16, +7, +11q, +12/12q	10–40 (GCB)
	+9/9p23–24, +2p13–16	50–90 (PMBCL)
B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and BL	t(8q24)/MYC	
	t(14;18)/IGH–BCL2	
	t(3q27)/BCL6	
Burkitt lymphoma	t(8;14)(q24;q32) and variants	100
	dup(1q)	~30–50
	+7, +12	~10–30

Plasma cell myeloma/ plasmacytoma	t(4;14)(p16;q32), t(11;14)(q13;q32), t(14;16)(q32;q23)	~20
	t(6;14)(p25;q32), t(8;14)(q24;q32)	5
	Other t(14q32)	
	Total t(14q32)	~40–60
	–13, del(13q)	15–50
	dup(1q), del(6q), del(11q)	5–20

transformation of FL to DLBCL has led to the suggestion that perhaps those diffuse lymphomas that carry a t(14;18) are the ones that have evolved from follicular NHL (Kneba et al., 1991).

The t(14;18)(q32;q21) in FL juxtaposes the *BCL2* oncogene next to the *IGH* locus (Tsujimoto et al., 1984). In molecular terms, the translocation is therefore different from the cytogenetically identical aberration involving *MALT1* in MALT lymphomas. In FL, the 18q21 breaks in 60–70% of the cases cluster within a 2.8 kb major breakpoint region (MBR) located in an untranslated part of the third and last (3') exon of *BCL2*; most of these breaks occur within much smaller subregions of the MBR (reviewed by Willis and Dyer, 2000; van Dongen et al., 2003). Some 10% of the breakpoints are found in the more distal minor cluster region (MCR). Variant breakpoints have also been described, including 3' *BCL2* (12%)



**Figure 2.2** | Translocations t(14;18)(q32;q21) and t(11;14)(q13;q32). Fluorescence R-banding. (A) Ideogram and partial karyotype with t(14;18)(q32;q21) involving *IGH* and *BCL2*. (B) Ideogram and partial karyotype with t(11;14)(q13;q32) involving *IGH* and *CCND1*.



and 5' MCR (6%) (Buchonnet et al., 2000, 2002). The breakpoints in the *IGH* locus fall within the region containing the *J*-segments. Indeed, the translocation is supposed to originate from a failed *VDJ* rearrangement, which in turn argues that this lymphoma derives from a precursor cell in the bone marrow rather than from a germinalcenter cell. Remarkably, the t(14;18)-positive clone seems to be characterized by considerable plasticity allowing the tumor cells to become microvascular endothelial cells (Streubel et al., 2004). The *BCL2* gene deregulated by the translocation encodes an inhibitor of apoptosis (for reviews, see Thomadaki and Scorilas, 2006; Zinkel et al., 2006; Letai, 2008). A subset of t(14;18)-positive lymphomas do not express intact *BCL2* due to somatic mutations of the gene (Schraders et al., 2005; Johnson et al., 2009; Adam et al., 2013). Vice versa, *BCL2* expression is not restricted to tumor cells of FL but can be observed in a wide range of B-cell lymphoid neoplasms including CLL and DLBCL.

Less common variants of t(14;18) are the *t(2;18) (p12;q21)* and *t(18;22)(q21;q11)* that juxtapose the *BCL2* gene next to the *IGK* and *IGL* loci, respectively (Lin et al., 2008). The breakpoints in these variants are located in the 5' end of the *BCL2* locus and not in the 3' end that typically is involved in t(14;18) (Adachi et al., 1990; Hillion et al., 1991). Like t(14;18), both these translocations also occasionally occur in CLL, mostly secondarily to trisomy 12 (Leroux et al., 1991; Bentley et al., 2005). Indeed, they might even be more common in that disease (Adachi et al., 1990; Lin et al., 2008). Due to their rarity, it is not yet proven whether FL with these variants show the same clinical course as do t(14;18)-positive FL.

Only in 10% of t(14;18)-positive cases is this the sole cytogenetic change (Johansson et al., 1995). Recurrent secondary alterations seen in 10% or more are +X, +1q21–44, +7, +12q, +18q, del(1)(p36), del(6q), del(10)(q22–24), and the development of polyploidy. The most frequent secondary event arising after the t(14;18) seems to be the duplication of the der(18)t(14;18) (Johansson et al., 1995; Horsman et al., 2001; Höglund et al., 2004; Aamot et al., 2007). The secondary chromosomal changes in t(14;18)-positive FL arise nonrandomly and in an apparently distinct temporal order. Four possible cytogenetic pathways have been shown to characterize the early stages of clonal evolution, which converge on a common route during later stages. Based on these pathways, FL with t(14;18) may be classified into cytogenetic subgroups defined by the presence or absence of 6q–, +7, or + der(18)t(14;18). These subgroups do not seem to be associated with distinctive prognostic features, however (Höglund et al., 2004).

The most prominent changes in t(14;18)-negative FL are numerical and structural changes of chromosome 3. Horsman et al. (2003) showed the average number of chromosomal aberrations to be similar in t(14;18)-positive and t(14;18)-negative FL, namely, 7.9 and 8.2, respectively. *t(14q32)* other than t(14;18) is present in 28% of t(14;18)-negative FL. *Rearrangements of 1p36* are significantly less frequent than in t(14;18)-positive FL (8% versus 18%), whereas *trisomy 3* is significantly more frequent (17% vs. 2%). Trisomy 3 may be part of a distinct cytogenetic pattern in t(14;18)-negative FL, which also includes *trisomy 18*. As described earlier, trisomies for chromosomes 3 and 18 are quite common in marginal zone lymphomas, and it must be questioned whether some of the +3 and +18 single- or double-positive lymphomas are not indeed marginal zone lymphomas with a follicular growth pattern rather than true FL.

In line with the higher frequency of FL grade 3 in t(14;18)-negative FL, several studies have documented *t(3q27)* to be highly recurrent in this disease. Horsman et al. (2003) found that *t(3;14) (q27;q32)* was significantly more frequent in lymphomas without than with t(14;18). The overall frequency of t(3q27) has been reported to be below 10% in t(14;18)-positive FL and FL of grades 1 and 2 but to reach 55% in FL3b with a DLBCL (Horsman et al., 2003; Katzenberger et al., 2004; Horn et al., 2011). In contrast to t(14;18)-positive FL, t(3q27)-positive FL less frequently express CD10 and *BCL2* (Jardin et al., 2002).

The target of t(3q27) is *BCL6*, which encodes a transcriptional regulator involved in germinal center formation (Jardin et al., 2007). The *BCL6* (formerly also called *BCL5* or *LAZ3*) gene was identified through cloning of the recurrent t(3;14) (q27;q32) involving *IGH* and was shown to be involved also in its light-chain variants *t(2;3) (p12;q27)* and *t(3;22)(q27;q11)* (Kerckaert et al., 1993; Ye et al., 1993a, 1993b; Miki et al., 1994a, 1994b). A steadily growing number of other t(3q27) translocations and intrachromosomal changes in chromosome 3 are also being cytogenetically identified and molecularly characterized, all of which juxtapose *BCL6* next to various partners (Table 2.2); for referencing and steady updates, the reader is referred to <http://atlasgeneticsoncology.org/Genes/BCL6ID20.html>. Indeed, *BCL6* seems to be one of the most promiscuous loci in B-cell lymphomas, and translocations affecting 3q27 are highly recurrent in various subtypes of B-cell disorders including particularly DLBCL or nodular lymphocyte predominance HL (discussed subsequently). The common theme of all these translocations is that the *BCL6* promoter is substituted by the promoter of the partner that interrupts its negative autoregulation, a similar effect to that re-

ported for somatic *BCL6* mutations (Chen et al., 1998; Pasqualucci et al., 2003). Despite this supposed common mechanism of action, considerable molecular heterogeneity of the breakpoints has been observed. First, an MBR in the 5' region of the gene is distinguished from the alternative breakpoint region (ABR) that is located more than 200 kb telomeric (Butler et al., 2002). With regard to the latter, it has not yet been ruled out that also other genes or regulatory elements than *BCL6* may be involved. Remarkably, *BCL6* breakpoints in t(14;18)-negative FL seem to predominantly affect the ABR, whereas those in DLBCL mostly affect the MBR (Bosga-Bouwer et al., 2005; Gu et al., 2009). Though the vast majority of t(3;14)(q27;q32) affects the constant regions of the *IGH* locus, switch gamma translocations seem to be more prominent in FL (70%), whereas switch  $\mu$  translocations are more frequent in DLBCL (89%) (Ruminy et al., 2006).

A distinctive subtype of t(14;18)-negative FL is characterized by a predominantly diffuse growth pattern and deletions in 1p36 (Katzenberger et al., 2008, 2009).

#### Clinical correlations and disease progression in FL

The early steps of clonal evolution and disease progression during lymphomagenesis have to be distinguished from the secondary changes that take place when already manifest FL transform to either higher-grade FL or high-grade NHL (e.g., DLBCL or DLBCL–BL intermediate). Several features are notable regarding the initial steps of FL lymphomagenesis:

1. The t(14;18)(q32;q21) can, in up to two-thirds of healthy individuals, be detected in nonmalignant B cells in the peripheral blood. Multiple clones may be present and can persist and expand over time (Limpens et al., 1995; Roulland et al., 2006a, 2014). The t(14;18) can also be found in reactive lymphoid tissue of healthy individuals (Limpens et al., 1991). At present, these cells may best be described as “follicular lymphoma-like cells” (FLLCs) (Roulland et al., 2006b).
2. The t(14;18)-positive FLLCs (and possibly also follicular lymphoma *in situ* (FLIS) cells; see following text) probably are able to reenter the germinal center and undergo additional rounds of somatic hypermutation there, after which they may circulate between the germinal center and bone marrow (Roulland et al., 2006b; Kluin, 2013). The recently identified t(14;18)-positive germinal center B cells represent an important missing link between the FLLCs and FLIS (see following text) (Kluin, 2014; Tellier et al., 2014).

3. So-called FLIS has been observed in lymph nodes from healthy individuals (Cong et al., 2002). These FLIS show strong *BCL2* expression but preservation of the nodal architecture. During the progression from FLIS to manifest FL, an increase in (epi)genomic complexity has been observed (Mamessier et al., 2014; Schmidt et al., 2014).

A model of nonpediatric FL has emerged, suggesting at least two cytogenetic pathways during the progression of manifest lymphomas:

FL grades 1 and 2 with t(14;18) can progress to FL3a when secondary changes occur or transform to secondary DLBCL through the acquisition of tertiary aberrations of, for example, 17p (*TP53*), 9p21 (*CDKN2A*), or 8q24 (*MYC*) (Christie et al., 2008). Transformation is mostly associated with an unfavorable prognosis (Johnson et al., 2008).

FL, mostly of grade 3b and lacking t(14;18) but frequently showing t(3q27), can also progress to DLBCL through pathways similar to those of the first group (Bosga-Bouwer et al., 2005, 2006). As for the (cytogenetic) pattern of clonal evolution, two main pathways can be distinguished: One includes linear clonal evolution in which the relapse arises from the previous FL clone (with the acquisition of additional genetic aberrations). The other involves divergent clonal evolution where the relapse originates from an FL precursor common to the tumor present at relapse and the preceding FL (Höglund et al., 2004; Berglund et al., 2007; Johnson et al., 2008; Green et al., 2013; Okosun et al., 2014; Pasqualucci et al., 2014; Loeffler et al., 2015). In addition to t(14;18), mutations of *CREBBP* and *MLL2* seem to be early events in FL lymphomagenesis.

A variety of cytogenetic and molecular cytogenetic studies have correlated chromosomal changes in FL with outcome. Tilly et al. (1994) showed that structural alterations of 1p21–22, 6q23–26, 4q, and 17p were significantly associated with reduced 5-year survival in FL. In t(14;18)-positive FL, del(1)(p36), del(6q), del(10)(q22–24), +7, the total number of abnormalities, the number of markers and additions, and the presence of polyploidy have been correlated with morphologic progression. In t(14;18)-positive FL, Höglund et al. (2004) reported +X, 1p–, +1q, +12, 17p–, and 17q– to be correlated with an unfavorable outcome in univariate analysis and +12 and del(17p) also correlated with poor survival in a multivariate analysis that included clinical risk factors. In male patients, gain of the X chromosome (as assessed by karyotyping and/or array CGH) correlated

with poor outcome (Johnson et al., 2008; Eide et al., 2010; Bouska et al., 2014). A cCGH study identified loss of 6q25–27 as being associated with unfavorable prognosis (Viardot et al., 2002), and recent aCGH-based reanalyses confirmed these findings and also suggested that deletions of 9p21 and gain of 11q are unfavorable prognostic markers (Schwaenen et al., 2009). In addition, high-resolution analysis has identified common sites of copy number-neutral LOH including 1p36 (with concomitant somatic mutations in *TNFRSF14*), 6p, 12q, and 16p (Cheung et al., 2010a, 2010b, 2012). *TNFRSF14* and *TP53* mutations have been associated with a worse prognosis (O'Shea et al., 2008; Cheung et al., 2010b). Gain/amplification of 2p15–16.1 (including the *REL* oncogene) may play a role in the transformation of FL to DLBCL (Kwiecinska et al., 2014). Several recent whole-genome and whole-exome sequencing studies have indicated an important role for chromatin modifiers, including *MLL2*, *CREBBP*, and *EZH2*, as driver and accelerator mutations in FL transformation (Green et al., 2013; Okosun et al., 2014; Pasqualucci et al., 2014). tDLBCL are more related to GCB-DLBCL than to ABC-DLBCL (Kwiecinska et al., 2014; Pasqualucci et al., 2014). All these studies should be regarded with some caution, however, since they were all retrospective and heterogeneous with regard to both treatment and t(14;18) positivity. Studies of large series of randomized FL patients in prospective trials are still lacking as are studies taking into account the gene expression risk profiles characteristic of this disease (Dave et al., 2004).

### Diffuse large B-cell lymphoma (DLBCL)

DLBCL are a morphologically, biologically, and clinically heterogeneous group that constitutes 30–40% of all B-cell lymphomas. Based on the site of involvement, some entities such as primary DLBCL of the CNS and leg-type primary cutaneous DLBCL are distinguished from systemic DLBCL (Jaffe et al., 2001; Swerdlow et al., 2008). Hardly any chromosomal data exist on these site-specific diseases, but interphase cytogenetic studies have indicated a similar spectrum of changes to that of systemic DLBCL (Hallermann et al., 2004a, 2004b).

Some quantitative differences may nevertheless exist, as more than 30% of leg-type DLBCL have been shown by FISH to have t(8;14)(q24;q32) or variants and also in the plasmablastic and immunoblastic variant was t(8q24) often seen by interphase cytogenetics (Hallermann et al., 2004a; Kanungo et al., 2006; Beaud-Rotureau et al., 2008; Aukema et al., 2014; Siebert et al., unpublished results). In addition, deletions of 6p21 involving the MHC region are reported in over 50% of lymphomas of immune-privileged sites (primary CNS and testicular

lymphomas) and are more common compared to nodal DLBCL (Riemersma et al., 2000; Booman et al., 2006, 2008; Schwindt et al., 2009). In primary lymphoma of the bone, gains of 1q and amplification of 2p16.1, both associated with a germinal center-like genetic profile, have been identified (Heyning et al., 2010).

Another subtype, ALK-positive DLBCL, in which the tumor cells show a typical immunoblastic morphology with expression of VS38 and IgA, was recognized based on the expression of anaplastic lymphoma kinase. These lymphomas are cytogenetically characterized by a simple or complex t(2;17)(p23;q23) involving the clathrin (*CLTC*) gene in 17q23 and the *ALK* gene in 2p23 (Chikatsu et al., 2003; Gascoyne et al., 2003; Gesk et al., 2005). They typically lack *MYC* rearrangements but instead show *MYC* gain and amplification (Valera et al., 2013).

Cytogenetic data on T-cell/histiocyte-rich large B-cell lymphoma (THRLBCL), an additional morphologic subtype of DLBCL, are scarce. CGH analysis of 17 THRLBCL showed genomic imbalances in all cases. The most common imbalances were gain of Xq, 4q13–28, Xp11–21, and 18q21 and loss of 17p (Franke et al., 2002).

The nodal and extranodal types of DLBCL mostly show centroblastic or immunoblastic morphology (Jaffe et al., 2001). These common DLBCL must be distinguished from primary mediastinal (thymic) large B-cell lymphoma (PMBCL) (Jaffe et al., 2001). Gene expression studies have shown the latter to be associated with a distinct expression signature related to that of cHL (Rosenwald et al., 2003; Savage et al., 2003). Various expression subgroups of DLBCL were also identified by these studies. Clinically, most relevant were two subgroups of DLBCL identified by gene expression profiling named according to the COO signature, that is, ABC-DLBCL associated with an unfavorable and GCB-DLBCL associated with a more favorable outcome (Alizadeh et al., 2000; Rosenwald et al., 2002; Gutierrez-García et al., 2011).

No chromosomal aberration is completely specific for DLBCL, but many of the generally lymphoma-associated rearrangements are prominent also in this disease subtype. t(3q27), involving the *BCL6* locus and various partners in a manner similar to that in FL, is seen in 20–40% of DLBCL (Table 2.2). Iqbal et al. (2007) detected *BCL6* translocations at the MBR by FISH in 25 (19%) of 133 DLBCL cases, with a higher frequency in PMBCL (33%) and ABC-DLBCL (24%) than in GCB-DLBCL (10%).

Translocations of 14q32 involving the *IGH* locus can be detected in up to half of all DLBCL (Lenz et al., 2007; Bouamar et al., 2013). Approximately 20–30% of



DLBCL show a t(14;18)(q32;q21) (Rosenwald et al., 2002; Hummel et al., 2006; Barrans et al., 2010; Visco et al., 2013). These cases are predominantly centroblastic and belong almost exclusively to the GCB-DLBCL subtype (Schlegelberger et al., 1999; Rosenwald et al., 2002; Iqbal et al., 2004; Visco et al., 2013). A t(8;14)(q24;q32) or variant t(8q24) can be detected in around 10% of DLBCL, some of which, especially (pediatric) cases with an *IG-MYC* translocation and lacking *BCL2* and *BCL6* rearrangements, might represent true BL based on their gene expression profile (Dave et al., 2006; Hummel et al., 2006; Klapper et al., 2008a; Aukema et al., 2014). In others, these translocations are supposed to be secondary changes mostly associated with a very aggressive clinical behavior and in part related to transformation into lymphomas intermediate between DLBCL and BL (see following text). In addition to t(3;14)(q27;q32) and the light-chain variants t(2;3)(p12;q27) and t(3;22)(q27;q11) involving *BCL6* and the *MYC*-targeting translocations more typical of BL, a steadily increasing number of other t(14q32) with various partners also exist (Lenz et al., 2007; Bouamar et al., 2013), particularly in the ABC-DLBCL tumor subset (Lenz et al., 2007).

Frequently recurring numerical chromosome aberrations in DLBCL include gains of chromosomes X, 3, 7, 12, and 18 and losses of the Y chromosome and chromosomes 6, 13, 15, and 17. Based on cytogenetic data, commonly gained chromosomal regions are 1q23–31, 3q21–22, 6p, 7p, 7q31–32, 8q22–24, 11q12–13, 12q14–24, and 18q11–21 in centroblastic lymphomas and 1q21–25, 3p24–21, 6p21, 7p12–21, 18q, and 22q12–qter in immunoblastic lymphomas. Commonly deleted chromosomal regions are 1p35–pter, 2p23–pter, 6q21–22, 6q25–qter, 8p12–pter, 9p21–pter, 11q23–qter, 12p12–13, and 17p12–13 in centroblastic lymphomas and 1p35–36, 2q22–24, 4q32–qter, 6q21–25, 7q33, 8q21, 9p24, 9q21–32, 11q21–qter, 14q23–qter, 16p13, and 18q21–qter in immunoblastic lymphomas. These imbalances mostly occur as part of complex karyotypes with many unbalanced translocations. Losses of the whole chromosome 10, deletions in 8q and 14q, as well as structural abnormalities of 4q have been reported to be significantly more frequent in immunoblastic than in centroblastic lymphomas (Johansson et al., 1995; Schlegelberger et al., 1999).

CGH studies suggest that the main effect of the vast majority of cytogenetic changes in DLBCL is to create a set of common imbalances that may, at least in part, form the basis for the subgroups of DLBCL defined by gene expression. ABC-DLBCL frequently show trisomy 3, gains of 3q (including *BCL6*), 18q21–22 (including *MALT1* and mostly *BCL2*), and 19q13 (including *SPIB*), but losses of

9p21 (including *CDKN2A*) and 6q21–22. GCBDLBCL show frequent gains of 1q, 2p13–16 (including *REL/BCL11A*), chromosome 7, 11q, and 12q12, whereas PMBCL show gains of 9p21–pter (including *JAK2/PDL2*) and 2p14–16 (including *REL/BCL11A*) (Bea et al., 2005; Tagawa et al., 2005; Kimm et al., 2007; Wessendorf et al., 2007; Lenz et al., 2008; Green et al., 2014). It should be stressed that complete or partial trisomy 9 with a minimal region of gain in 9p23–24 seems to be specific for PMBCL (Bentz et al., 2001). In general, specific chromosomal alterations might be associated with significant changes in gene expression signatures that reflect various aspects of lymphoma cell biology as well as the host response to the lymphoma (Bea et al., 2005).

Next-generation sequencing analyses have revealed a large number of potentially targetable, recurrently mutated genes including *MLL2*, *TP53*, *TNFRSF14*, *CARD11*, *GNA13*, *CD79B*, *EZH2*, *MYD88*, and *CREBPP* (Ngo et al., 2011; Pasqualucci et al., 2011a, 2011b; Lohr et al., 2012; Morin et al., 2013; Zhang et al., 2013; Bohers et al., 2014; Vaqué et al., 2014). Some of these are associated with one of the COO types, for example, *EZH2* (15–30%) and *GNA13* (15–30%) mutations in GCB-DLBCL and *MYD88* (20–35%) and *CD79B* (15–20%) mutations in ABC-DLBCL (Morin et al., 2011; Ngo et al., 2011; Pasqualucci et al., 2011a; Zhang et al., 2013; Bohers et al., 2014).

### Clinical correlations in DLBCL

All clinical correlations in DLBCL have to be treated with extreme caution since the data derive from before the introduction of anti-CD20 antibody therapy (Rituxan). Moreover, established clinical risk factors including International Prognostic Index (IPI) (The International Non-Hodgkin's Lymphoma Prognostic Factors Project, 1993), subtyping by gene expression profiling, and evaluation in the context of other chromosomal aberrations were mostly not taken into account. The data on the prognostic importance of t(14;18) (Barrans et al., 2003; Akyurek et al., 2012; Visco et al., 2013; Bellas et al., 2014) and t(3q27) (Akasaka et al., 2000b; Barrans et al., 2002; Ueda et al., 2002a; Niitsu et al., 2007; Shustik et al., 2010; Akyurek et al., 2012; Akay et al., 2014) are conflicting; at present, no convincing evidence exists that one or the other is reliably linked to prognosis. Like in many other lymphomas, there is good evidence that del(17p) and del(9)(p21) in DLBCL are associated with progression and unfavorable prognosis (Tagawa et al., 2005). In centroblastic DLBCL, deletions and duplications of 1q have been reported to be adverse risk factors independent of IPI, whereas trisomy 5 and changes of 15q seem to be independent indicators of a reduced risk. In immunoblastic lym-

phomas, changes of 7q and 8q had a stronger impact on survival than did the IPI (Schlegelberger et al., 1999). Gains of *MAPKAPK3* located at 3p21.31 and loss of 9p21.3 (*CDKN2A/B*) have been associated with chemoresistance in DLBCL patients treated with immunochemotherapy (Kreisel et al., 2011). Interestingly, another study using array CGH found that clonal heterogeneity and 9p21.3 loss were associated with poor prognosis (Suguro et al., 2014). In addition to genomic imbalances, also aberrant DNA methylation has been shown to have prognostic impact with heavily methylated DLBCL cases showing a worse clinical outcome (Chambwe et al., 2014).

Recent advances in gene expression profiling and assessment of cytogenetic imbalance patterns offer new ways to classify lymphomas into prognostically meaningful subgroups based on their COO (Masque-Soler et al., 2013). Gains of 3p11–12 provide prognostic information independently of the gene expression data (Bea et al., 2005). It is not likely, however, that this way of subgrouping patients exhausts the possibilities for clinically meaningful classification. DLBCL can show considerable molecular genetic heterogeneity, including aberrant somatic hypermutation of *IGH*, *BCL6*, *MYC*, and other genes (Pasqualucci et al., 2001). This and frequent uniparental disomy and epigenetic changes are biological parameters also likely to influence outcome. The adverse prognostic impact of *MYC* (in combination with additional *BCL2* and/or *BCL6*) rearrangements in DLBCL has been demonstrated in multiple FISH studies (Klapper et al., 2008b; Savage et al., 2009; Barrans et al., 2010; Aukema et al., 2014; Tzankov et al., 2014). It is therefore advisable to screen all DLBCL in routine diagnostic situations for *MYC* and related breakpoints with both break-apart and dualcolor dual fusion probes, in addition to performing chromosome banding analysis with karyotyping (Klapper et al., 2008b; Savage et al., 2009; Barrans et al., 2010; May et al., 2010; Kluin and Schuurin, 2011; Aukema et al., 2011, 2014; Tzankov et al., 2014). In the near future, these molecular cytogenetic analyses will likely be combined with targeted panel sequencing for diagnostic and prognostic relevant genes and COO classification using dedicated gene expression profiling.

### Other large B-cell lymphomas

Other very rare entities of large B-cell lymphoma include intravascular large B-cell lymphoma (ILBL) and primary effusion lymphoma (PEL) (Jaffe et al., 2001; Swerdlow et al., 2008). In the former disease, hardly any cytogenetic studies have been reported though the presence of t(11;14)(q13;q32), t(14;18)(q32;q21), and *BCL6* rearrangements has been described in single cases (Vieites et al.,

2005; Rashid et al., 2006; Cui et al., 2014). A recent review by Mandal et al. (2007) suggested that structural aberrations of chromosomes 1, 6, and 10 and, less frequently, chromosomes 4, 5, and 8 might be nonrandom changes with a recurrent site of deletion in 6q21–24. PEL is universally associated with human herpes virus 8 (HHV-8)/Kaposi sarcoma herpesvirus (KSHV) infection. Also, EBV is found in most cases. Cytogenetic analyses mostly show a complex karyotype with structural rearrangements frequently involved in lymphomas in general but without subgroup-specific abnormalities (Boulanger et al., 2001). Trisomy 7, trisomy 12, and aberrations proximally in the long arm of chromosome 1 seem to be recurrent (Wilson et al., 2002). A CGH study identified recurrent gains of the X chromosome and 12q (Mullaney et al., 2000). In another array CGH study, copy number abnormalities involving, among others, 1q, 7q, 8q, 14q32, and chromosome 12 were identified with *SELPLG/CORO1C* at 12q24.1 showing amplification/lowlevel gain in 23% of the cases (Luan et al., 2010).

The majority of PEL occur in the setting of HIV infection, which is associated with an increased risk of lymphoma. AIDS-related lymphomas include systemic lymphomas, mostly DLBCL and BL, primary central nervous system lymphomas (PCNSL), PEL, and plasmablastic lymphoma of the oral cavity (Jaffe et al., 2001; Swerdlow et al., 2008). AIDS-related and other lymphomas in immunodeficient patients show a spectrum of chromosomal aberrations similar to that of their counterparts in immunocompetent patients (Vaghefi et al., 2006). Activation of *MYC* and inactivation of *TP53* in conjunction with EBV and other viral infections seem to play a major pathogenic role (Carbone, 2003). In line with these molecular changes, there seems to be a trend toward more complex karyotypes and a higher frequency of t(8;14)(q24;q32) and the variant translocations t(2;8) and t(8;22). In lymphomatoid granulomatosis, which is also an Epstein–Barr virus (EBV)-driven lymphoproliferative disorder for which immunodeficient patients carry an increased risk, no cytogenetic aberrations have yet been identified in line with the oligo- or polyclonal origin documented particularly in low-grade cases (Jaffe et al., 2001; Swerdlow et al., 2008).

### Burkitt lymphoma (BL)

BL is a highly aggressive disease often presenting at extranodal sites or as an acute leukemia. It is the most common B-cell lymphoma in the pediatric age group but is also observed in the adult population (Boerma et al., 2004; Leoncini et al., 2008). Endemic and sporadic BL are recognized as variants manifesting differences in clinical presentation and biology. Immunodeficiency-associated BL is seen primarily in association with HIV infection.




The cytogenetic hallmark of BL is the t(8;14) (q24;q32) and its variants, t(2;8) (p12;q24) and t(8;22)(q24;q11). These translocations are present both in the endemic, African tumor type and in the sporadic BL occurring in Europe, America, and Japan, both in EBV-infected patients and in EBVnegative BL. The t(8;14) is present in 75–85% of all BL (Johansson et al., 1995). In 15–25%, one of the variant translocations is found with the t(8;22) occurring twice as frequently as t(2;8) (Figure 2.3).

BL was the first lymphoid neoplasm in which the underlying chromosomal rearrangement was characterized. The first cytogenetic study of BL was reported in 1963 by Jacobs and coworkers. After the introduction of banding techniques, Manolov and Manolova (1972) described an additional band at the end of 14q in five of six fresh BL and in seven of nine cell lines from such tumors. The nature of the BL-specific abnormality was clarified by Zech et al. (1976), who established that the 14q+ arose through the translocation t(8;14)(q24;q32). Based on a high-resolution study, Manolova et al. (1979) reported that the translocation was reciprocal. Further scrutiny of prophase–prometaphase cells from the BL-derived cell line Daudi allowed Zhang et al. (1982) to map the translocation breakpoints to subbands 8q24.13 and 14q32.33. Berger et al. (1979) described the first variant translocation, t(8;22)(q24;q11), and Miyoshi et al. (1979) and van den Berghe et al. (1979) simultaneously reported the second variant, t(2;8)(p12;q24).

The molecular consequence of the t(8;14) and its variants is deregulation of the *MYC* oncogene in 8q24 through juxtaposition next to one of the *IG* enhancer elements in the *IGH* (14q32), *IGK* (2p12), or *IGL* (22q11) locus. Activation of *MYC* takes place on the der(14) in t(8;14) and on the der(8) in t(2;8) and t(8;22). Correspondingly, the breakpoints on the der(8) differ between the t(8;14) and its variants being telomeric of the *MYC* gene in the latter and centromeric of *MYC* in the former. Breakpoints can scatter over several hundred kilobases on both sides of the *MYC* locus (Joos et al., 1992a, 1992b; Einerson et al., 2006; Kroenlein et al., 2012). Sporadic, endemic, and immunodeficiency-associated BL show different clustering of the breakpoints on der(14) and der(8), indicating that different pathogenetic mechanisms may be behind the generation of the t(8;14) in these different disease settings. Briefly, the breakpoints of t(8;14) in endemic BL frequently fall far centromeric of the *MYC* gene and regularly affect the J-segments of the *IGH* locus. In contrast, t(8;14) in sporadic BL more frequently involves the switch regions of the constant segments of *IGH*, or the breakpoint occurs between exons 1 and 2 or immediately 5' of the *MYC* gene. This molecular hetero-

geneity has to be taken into account when diagnosing one of the translocations by molecular or molecular cytogenetic techniques, and therefore, it is advisable always to use both a *MYC* break-apart and *MYC-IGH* fusion FISH assay as far



**Figure 2.3** | Burkitt translocation t(8;14) and variants. (A) Ideogram and partial karyotype with t(8;14)(q24;q32) involving *MYC* and *IGH*. (B) Ideogram and partial karyotype with t(2;8)(p12;q24) involving *IGK* and *MYC*. (C) Ideogram and partial karyotype with t(8;22)(q24;q11) involving *MYC* and *IGL*. Fluorescence R-banding.

centromeric breakpoints may be missed using a break-apart assay alone (Hummel et al. 2006). In addition, somatic mutation of the *MYC* locus occurs frequently in BL, which might in part be driven by the VH mutation machinery after the juxtaposition of *MYC* next to *IGH*. Some of the mutations in *MYC*, which may also occur independently of *IGH*-driven hypermutation, might be of pathogenetic relevance. For more details on the molecular characteristics of the Burkitt translocations as well as the pathogenic consequences of *MYC* activation in general, the reader is referred to excellent reviews by, for example, Willis and Dyer (2000), Boxer and Dang (2001), Küppers and Dalla-Favera (2001), Küppers (2005), Klapproth and Wirth (2010), and Ott et al. (2013).

Typically in BL, the t(8;14) or variant translocation is part of a rather simple karyotype; complex karyotypes indicate disease progression and are an adverse prog-

nostic factor (Johansson et al., 1995; Hummel et al., 2006; Salaverria et al., 2008; Boerma et al., 2009; Poirel et al., 2009). Secondary chromosomal changes are present in around 60% of BL (Berger et al., 1989; Johansson et al., 1995; Boerma et al., 2009). In a comparison of paired karyotypes from BL patients, complex karyotypes were seen at a higher frequency in relapsed or progressive BL compared to initial diagnostic samples (Aukema and Siebert, in prep.); this may contribute to the very dismal prognosis of patients with progressing and relapsing disease (Woessmann, 2013). The clearly most frequent secondary aberration (>30% of all patients) is structural rearrangement of chromosome 1, in particular of the long arm and often leading to partial trisomy for 1q. Trisomy 7 and trisomy 12 are other common secondary changes. These three aberrations tend to be mutually exclusive in BL progression (Boerma et al., 2009). Another hot spot for secondary involvement in BL is chromosomal region 13q3, found rearranged in 15% (Johansson et al., 1995). The target is most likely the *miR-17-92* miRNA locus that, like other recurrent molecular changes, seems to interact with *MYC* in transformation (He et al., 2005; O'Donnell et al., 2005). Losses of 13q may be associated with a more unfavorable outcome (Nelson et al., 2010). In addition to secondary chromosomal changes, whole-genome and whole-exome sequencing studies have identified recurrent mutations in *MYC*, *TP53*, *ID3*, *TCF3*, *CCND3*, and *DDX3X* as common, with mutations targeting *ID3* or *TCF3* occurring in about 70% of sporadic BL cases (Campo, 2012; Love et al., 2012; Richter et al., 2012; Schmitz et al., 2012).

Although the putative existence of BLs lacking a *MYC* translocation has been subject of much discussion (Leucci et al., 2008), it is unlikely that “true” *MYC* translocation-negative BL exist. Recently, in a subset of lymphomas resembling BL by gene expression profile and pathological characteristics but importantly lacking *MYC* translocation, 11q aberrations with proximal gains and telomeric losses were identified (Salaverria et al., 2014).

The view that BL is a homogeneous disease characterized by the presence of one of the three specific translocations is also supported by gene expression profiling studies (Dave et al., 2006; Hummel et al., 2006). Nevertheless, none of the three translocations is completely specific for BL as all of them have been described also in several other B-cell lymphomas. “B-cell lymphoma with features intermediate between DLBCL and BL” is a disease subset strongly enriched for the presence of these translocations and might even present with a typical Burkitt leukemia morphology in the bone marrow (Kluin et al., 2008). In contrast to BL, these intermediate lymphomas, which occur at a significantly older age,

mostly show complex karyotypes (Hummel et al., 2006; Boerma et al., 2009; Salaverria and Siebert, 2011; Aukema et al., 2014). Many of them harbor other recurrent *IG* translocations such as t(14;18)(q21;q32) or t(3q27) targeting the *BCL6* locus. These so-called “doublehit” lymphomas are associated with a much more unfavorable outcome, highlighting the clinical relevance of distinguishing both subtypes (Dave et al., 2006; Hummel et al., 2006; Aukema et al., 2011). Moreover, a wide range of t(8q24) that do not involve *IG* loci as partners, such as t(8;9)(q24;p13) juxtaposing *MYC* to *PAX5* and t(3;8)(q27;q24) juxtaposing *MYC* to *BCL6* (Hummel et al., 2006; Bertrand et al., 2007; Sonoki et al., 2007), may also occur. The detection of a double hit, the presence of a non-*IG*-*MYC* translocation, and the occurrence of a t(8;14) or variant as part of a complex karyotype should, particularly in nonpediatric patients, lead to a reassessment of the histopathologic diagnosis, especially since each of these three features is associated with an unfavorable outcome (Hummel et al., 2006; Boerma et al., 2009; Aukema et al., 2011, 2014; Salaverria and Siebert, 2011). In cases where a diagnosis of BL is in doubt, profiling for BL-specific genes (the analysis can also be performed on formalin-fixed and paraffin-embedded tissues) as well as mutational analysis (e.g., for *ID3* and *TCF3* mutations that are absent in other *MYC* translocation-positive lymphomas) may be helpful in establishing the correct diagnosis (Love et al., 2012; Richter et al., 2012; Schmitz et al., 2012; Masque-Soler et al., 2013).

### Clinical correlations

The cytogenetic changes in mature lymphoid neoplasms and their clinical impact greatly depend on the disease subtype in which they occur. They have therefore mostly been discussed in the respective disease-specific sections.

The most important clinical reason for identifying chromosomal aberrations in suspected mature lymphoid neoplasms is the fact that they provide essential information on which a correct, precise diagnosis can be based. Different histopathologic subtypes of mature lymphoid neoplasms are associated with very different natural courses and show quite different responses to therapy. Modern therapy for mature lymphoid neoplasms relies on the proper classification of the lymphoma, and the patterns of both primary and secondary chromosomal aberrations constitute part of the foundation for that classification. Indeed, comments on the 2001 WHO classification recommend that cytogenetic analysis be performed in all cases and should form part of the pathology report (Harris et al., 1999, 2000).

It has to be emphasized that the same chromosomal aberrations may have different prognostic impact in the various subtypes of mature lymphoid neoplasms. Among NHL, for example, t(11;14) (q13;q32) mostly signifies a MCL that has a poor prognosis, although a subgroup of MCL with this translocation and favorable prognosis exists (Navarro et al., 2012). In contrast, t(11;14) in MM defines a disease subset with a relatively favorable outcome (though compared to t(11;14)-positive MCL, the overall survival of patients with t(11;14)-positive MM is still worse). Moreover, t(11;14) can be detected in preneoplastic disorders like MGUS and in quite indolent diseases like SLVL. These findings point to the need to integrate cytogenetic analysis into an interdisciplinary diagnostic workup of mature lymphoid neoplasms, which, besides histopathologic features and the immunophenotype, also takes into account both the clinical presentation and the neoplastic cells' genetic characteristics. It can be anticipated that the diagnostic process will soon also see the integration of findings made by array-based technologies like gene expression profiling as well as high-throughput sequencing.

Despite the fact that the clinical relevance of cytogenetic changes in mature lymphoid neoplasms always has to be evaluated in the context of the disease subtype, a few common themes generally associated with unfavorable outcome seem to exist: in most instances of mature lymphoid neoplasms, the rule of thumb is that the more complex the karyotype, the worse is the prognosis. A notable exception here is cHL that is characterized by highly complex karyotypes, and yet long-term cure rates far exceeding 80% are reached with modern therapy. Losses of 17p and 9p are supposed to mostly target the *TP53* and *TP16* gene loci. One or both changes are associated with unfavorable outcome in many mature lymphoid neoplasms, including CLL, FL, MCL, and DLBCL. Finally, translocations affecting the band 8q24 mostly target the *MYC* oncogene. In most types of mature lymphoid neoplasms, including FL, MCL, and DLBCL, these changes are associated with increased clinical aggressiveness and an unfavorable prognosis. The notable exception here is BL in which t(8;14) and variants are the pathognomonic alterations. Despite being a biologically highly aggressive disease, BL can be cured in most instances, particularly in children.

The near future will see the increasing introduction of molecular cytogenetic techniques like FISH and array CGH into the diagnosis of mature lymphoid neoplasms. Indeed, interphase FISH is already widely used for the diagnosis of the most recurrent alterations in CLL and MM and is also becoming a routine tool for the detection of diagnostically relevant chromosomal alterations in for-

malin-fixed, paraffin-embedded tissues (Ventura et al., 2006). Nevertheless, in contrast to chromosome banding analysis, interphase FISH will never provide a full overview of all chromosomal aberrations of a mature lymphoid neoplasm. This carries the very real risk that clinically relevant alterations, like t(8q24) in t(14;18)-positive FL, might be missed simply because they were not searched for. Moreover, a subclone with prognostically unfavorable chromosomal aberrations may be missed by FISH, whereas it might be detected by conventional cytogenetics due to a proliferative advantage of these cells. Finally, complex and variant FISH patterns might be difficult to interpret without knowledge of the metaphase chromosomal picture, the karyotype.

The advent of array CGH and related techniques enables the production of a wealth of new data at a resolution somewhere between the cytogenetic and molecular levels. One should exercise considerable caution when trying to translate findings and conclusions between the microscopic cytogenetic and the array-based information realms. As can be seen from the previous discussion of 13q deletions in MM, the diagnostic impact of a chromosomal change may seemingly depend on the technique with which it was detected (Dewald et al., 2005). Both small and large genetic changes can be missed if one exclusively relies on only one technique, not least if the change in question is present only in a subclone. The results are invariably best if different techniques and investigative principles are allowed to complement each other.

Recent technological developments have allowed high-throughput sequencing to become a part of both scientific and diagnostic analyses of lymphatic neoplasms. These new techniques provide valuable information helping us to understand the pathogenesis of neoplastic diseases better as well as to make more meaningful diagnoses and choose the optimal treatment. Prominent examples are *ID3* mutations in BL, *MYD88* mutations in Waldenström disease, or *RHOA* mutations in ATLL. In principle, sequencing technologies are able to detect all kinds of genomic alterations, be they balanced translocations, imbalances, or mutations. In practice, this may not be so easy, not least when it comes to translocations with repeat-rich breakpoints. In addition, the subclonal composition of tumors is hard to evaluate if these techniques are used alone. Finally, a wealth of bioinformatic challenges need to be resolved for daily use. It nevertheless seems beyond doubt that the future will see an integration of the different tumor genetic techniques also for diagnostic purposes, in lymphatic neoplasms

as elsewhere, with important clinical benefits for the patients (Biesecker et al., 2012; Yates and Campbell, 2012; Mwenifumbo and Marra, 2013).

It should finally be stressed that the treatment of many mature lymphoid neoplasms is now undergoing profound changes. Probably, the most important recent improvement in the treatment of the most common lymphomas, including the mature B-cell lymphomas MCL, FL, and DLCL, was the introduction of anti-CD20 immunotherapy (Rituxan) (Coiffier et al, 2002; Kluin-Nelemans et al., 2012). Almost all cytogenetic studies referred to earlier were performed before the introduction of this therapy, and we do not know whether the cytogenetic–prognostic correlations detected during the pre-Rituxan era still apply when the new antibody treatment is given. Indeed, with the exception of CLL and MM and in part BL and DLCL, hardly any systematic cytogenetic studies within clinical trials exist. This needs to be changed in future so that cytogenetic studies as well as molecular studies form an essential and integral part of prospective clinical trials; only then can the clinical impact of the various acquired genomic changes in mature lymphoid neoplasms be reliably assessed.

### Acknowledgments

The authors' own studies on chromosomal aberrations in lymphoid neoplasms are supported by Bundesministerium für Bildung und Forschung (BMBF), Deutsche Krebshilfe, Wilhelm Sander-Stiftung, KinderKrebsInitiative (KKI) Buchholz/Holm-Seppensen, European Union, Lymphoma Research Foundation (New York), and Schleswig-Holsteinische Krebsgesellschaft. The authors would like to thank Dr. Jose-Ignacio Martin-Subero, Dr. Borja Saez, and Heike Blohm for editorial support and Dr. Lana Harder, Dr. Simone Heidemann, and Claus-Peter Blohm for providing the figures and the graphical design. Finally, the authors acknowledge all colleagues worldwide who have made publicly available their cytogenetic data on lymphoid neoplasms and at the same time apologize to all whose work could not be cited herein. SMA is a fellow of the JSM-UMCG MD-PhD program, a recipient of the "Nijbakker-Morra" and "Hippocrates" Foundations awards, and is supported by the "Foundation de Drie Lichten," the "René Vogels' Foundation," and the framework of a "JSM-Ubbo Emmius Foundation Talent Grant."

## REFERENCES

- Aamot HV, Bjørnslett M, Delabie J, Heim S (2005): t(14;22) (q32;q11) in non-Hodgkin lymphoma and myeloid leukaemia: molecular cytogenetic investigations. *Br J Haematol* 130:845–851.
- Aamot HV, Torlakovic EE, Eide MB, Holte H, Heim S (2007): Non-Hodgkin lymphoma with t(14;18): clonal evolution patterns and cytogenetic–pathologic–clinical correlations. *J Cancer Res Clin Oncol* 133:455–470.
- Abdul-Wahab A, Tang SY, Robson A, Morris S, Agar N, Wain EM, et al. (2014): Chromosomal anomalies in primary cutaneous follicle center cell lymphoma do not portend a poor prognosis. *J Am Acad Dermatol* 70:1010–1020.
- Adachi M, Tefferi A, Greipp PR, Kipps TJ, Tsujimoto Y (1990): Preferential linkage of bcl-2 to immunoglobulin light chain gene in chronic lymphocytic leukemia. *J Exp Med* 171:559–564.
- Adam P, Baumann R, Schmidt J, Bettio S, Weisel K, Bonzheim I, et al. (2013): The BCL2 E17 and SP66 antibodies discriminate 2 immunophenotypically and genetically distinct subgroups of conventionally BCL2-"negative" grade 1/2 follicular lymphomas. *Hum Pathol* 44:1817–1826.
- Agopian J, Navarro JM, Gac AC, Lecluse Y, Briand M, Grenot P, et al. (2009): Agricultural pesticide exposure and the molecular connection to lymphomagenesis. *J Exp Med* 206:1473–1483.
- Akagi T, Motegi M, Tamura A, Suzuki R, Hosokawa Y, Suzuki H, et al. (1999): A novel gene, MALT1 at 18q21, is involved in t(11;18) (q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene* 18:5785–5794.
- Akao Y, Seto M, Yamamoto K, Iida S, Nakazawa S, Inazawa J, et al. (1992): The RCK gene associated with t(11;14) translocation is distinct from the MLL/ALL-1 gene with t(4;11) and t(11;19) translocations. *Cancer Res* 52:6083–6087.
- Akasaka T, Miura I, Takahashi N, Akasaka H, Yonetani N, Ohno H, et al. (1997): A recurring translocation, t(3;6) (q27;p21), in non-Hodgkin's lymphoma results in replacement of the 5' regulatory region of BCL6 with a novel H4 histone gene. *Cancer Res* 57:7–12.
- Akasaka H, Akasaka T, Kurata M, Ueda C, Shimizu A, Uchiyama T, et al. (2000a): Molecular anatomy of BCL6 translocations revealed by long-distance polymerase chain reaction-based assays. *Cancer Res* 60:2335–2341.
- Akasaka T, Ueda C, Kurata M, Akasaka H, Yamabe H, Uchiyama T, et al. (2000b): Nonimmunoglobulin (non-Ig)/ BCL6 gene fusion in diffuse large B-cell lymphoma results in worse prognosis than Ig/BCL6. *Blood* 96:2907–2909.
- Akasaka T, Lossos IS, Levy R (2003): BCL6 gene translocation in follicular lymphoma: a harbinger of eventual transformation to diffuse aggressive lymphoma. *Blood* 102:1443–1448.
- Akay OM, Aras BD, Isiksoy S, Toprak C, Mutlu FS, Artan S, et al. (2014): BCL2, BCL6, IGH, TP53 and MYC protein expression and gene rearrangements as prognostic markers in diffuse large B-cell lymphoma: a study of 44 Turkish patients. *Cancer Genet* 207:87–93.
- Akyurek N, Uner A, Benekli M, Barista I (2012): Prognostic significance of MYC, BCL2, and BCL6 rearrangements in patients with diffuse large B-cell lymphoma treated with cyclophosphamide, doxorubicin, vincristine, and prednisone plus rituximab. *Cancer* 118:4173–183.
- Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. (2013): Signatures of mutational processes in human cancer. *Nature* 500:415–421.
- Alizadeh AA, Eisen MB, Davis RE, Ma C, Lossos IS, Rosenwald A, et al. (2000): Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 403:503–511.



- Ansell SM, Akasaka T, McPhail E, Manske M, Braggio E, Price-Troska T, et al. (2012): t(X;14) (p11;q32) in MALT lymphoma involving GPR34 reveals a role for GPR34 in tumor cell growth. *Blood* 120:3949–3957.
- Attarbaschi A, Beishuizen A, Mann G, Rosolen A, Mori T, Uyttebroeck A, et al. (2013): Children and adolescents with follicular lymphoma have an excellent prognosis with either limited chemotherapy or with a “Watch and wait” strategy after complete resection. *Ann Hematol* 92:1537–1541.
- Au WY, Gascoyne RD, Viswanatha DS, Skinnider BF, Connors JM, Klasa RJ, et al. (1999): Concurrent chromosomal alterations at 3q27, 8q24 and 18q21 in B-cell lymphomas. *Br J Haematol* 105:437–440.
- Au WY, Horsman DE, Viswanatha DS, Connors JM, Klasa RJ, Gascoyne RD (2000): 8q24 translocations in blastic transformation of mantle cell lymphoma. *Haematologica* 85:1225–1227.
- Au WY, Gascoyne RD, Viswanatha DS, Connors JM, Klasa RJ, Horsman DE (2002): Cytogenetic analysis in mantle cell lymphoma: a review of 214 cases. *Leuk Lymphoma* 43:783–791.
- Au WY, Horsman DE, Gascoyne RD, Viswanatha DS, Klasa RJ, Connors JM (2004): The spectrum of lymphoma with 8q24 aberrations: a clinical, pathological and cytogenetic study of 87 consecutive cases. *Leuk Lymphoma* 45:519–528.
- Aukema SM, Siebert R, Schuurin E, van Imhoff GW, Kluin-Nelemans HC, Boerma EJ, Kluin PM (2011): Double-hit B-cell lymphomas. *Blood* 117:2319–2331.
- Aukema SM, Kreuz M, Kohler CW, Rosolowski M, Hasenclever D, Hummel M, et al. (2014): Biological characterization of adult MYC-translocation-positive mature B-cell lymphomas other than molecular Burkitt lymphoma. *Haematologica* 99:726–735.
- Banks PM, Chan J, Cleary ML, Delsol G, DeWolf-Peters C, Gatter K, et al. (1992): Mantle cell lymphoma. A proposal for unification of morphologic, immunologic, and molecular data. *Am J Surg Pathol* 16:637–640.
- Barrans SL, O'Connor SJ, Evans PA, Davies FE, Owen RG, Haynes AP, et al. (2002): Rearrangement of the BCL6 locus at 3q27 is an independent poor prognostic factor in nodal diffuse large B-cell lymphoma. *Br J Haematol* 117:322–332.
- Barrans SL, Evans PA, O'Connor SJ, Kendall SJ, Owen RG, Haynes AP, et al. (2003): The t(14;18) is associated with germinal center-derived diffuse large B-cell lymphoma and is a strong predictor of outcome. *Clin Cancer Res* 9:2133–2139.
- Barrans S, Crouch S, Smith A, Turner K, Owen R, Patmore R, et al. (2010): Rearrangement of MYC is associated with poor prognosis in patients with diffuse large B-cell lymphoma treated in the era of rituximab. *J Clin Oncol* 28:3360–3365.
- Barth TF, Bentz M, Leithäuser F, Stilgenbauer S, Siebert R, Schlotter M, et al. (2001): Molecular-cytogenetic comparison of mucosa-associated marginal zone B-cell lymphoma and large B-cell lymphoma arising in the gastrointestinal tract. *Genes Chromosomes Cancer* 31:316–325.
- Bäsecke J, Griesinger F, Trümper L, Brittinger G (2002): Leukemia- and lymphoma-associated genetic aberrations in healthy individuals. *Ann Hematol* 81:64–75.
- Bastard C, Tilly H, Lenormand B, Bigorgne C, Boulet D, Kunlin A, et al. (1992): Translocations involving band 3q27 and Ig gene regions in non-Hodgkin's lymphoma. *Blood* 79:2527–2531.
- Bauer B, Siebert R, Traulsen A. (2014): Cancer initiation with epistatic interactions between driver and passenger mutations. *J Theor Biol* 358:52–60.
- Bea S, Zettl A, Wright G, Salaverria I, Jehn P, Moreno V, et al. (2005): Diffuse large B-cell lymphoma subgroups have distinct genetic profiles that influence tumor biology and improve gene-expression-based survival prediction. *Blood* 106:3183–3190.
- Beà S, Valdes-Mas R, Navarro A, Salaverria I, Martin-Garcia D, Jares P, et al. (2013): Landscape of somatic mutations and clonal evolution in mantle cell lymphoma. *Proc Natl Acad Sci USA* 110:18250–18255.
- Belaud-Rotureau MA, Marietta V, Vergier B, Mainhaguiet G, Turmo M, Idrissi Y, et al. (2008): Inactivation of p16INK4a/CDKN2A gene may be a diagnostic feature of large B cell lymphoma leg type among cutaneous B cell lymphomas. *Virchows Arch* 452:607–620.
- Bellas C, García D, Vicente Y, Kilany L, Abaira V, Navarro B, et al. (2014): Immunohistochemical and molecular characteristics with prognostic significance in diffuse large B-cell lymphoma. *PLoS One* 9:e98169.
- Bentley G, Palutke M, Mohamed AN (2005): Variant t(14;18) in malignant lymphoma: a report of seven cases. *Cancer Genet Cytogenet* 157:12–17.
- Bentz M, Barth TF, Brüderlein S, Bock D, Schwertner MJ, Baudis M, et al. (2001): Gain of chromosome arm 9p is characteristic of primary mediastinal B-cell lymphoma (MBL): comprehensive molecular cytogenetic analysis and presentation of a novel MBL cell line. *Genes Chromosomes Cancer* 30:393–401.
- Berger R, Bernheim A, Weh HJ, Flandrin G, Daniel MT, Brouet JC, et al. (1979): A new translocation in Burkitt's tumor cells. *Hum Genet* 53:111–112.
- Berger R, Le Coniat M, Derré J, Vecchione D (1989): Secondary nonrandom chromosomal abnormalities of band 13q34 in Burkitt lymphoma-leukemia. *Genes Chromosomes Cancer* 1:115–118.
- Bertrand P, Bastard C, Maingonnat C, Jardin F, Maisonneuve C, Courel MN, et al. (2007): Mapping of MYC breakpoints in 8q24 rearrangements involving non-immunoglobulin partners in B-cell lymphomas. *Leukemia* 21:515–523.
- Berglund M, Enblad G, Thunberg U, Amini RM, Sundström C, Roos G, et al. (2007): Genomic imbalances during transformation from follicular lymphoma to diffuse large B-cell lymphoma. *Mod Pathol* 20:63–75.
- Biagi JJ, Seymour JF (2002): Insights into the molecular pathogenesis of follicular lymphoma arising from analysis of geographic variation. *Blood* 99:4265–4275.
- Biesecker LG, Burke W, Kohane I, Plon SE, Zimmern R (2012): Next-generation sequencing in the clinic: are we ready? *Nat Rev Genet* 13:818–824.
- Boerma EG, van Imhoff GW, Appel IM, Veeger NJ, Kluin PM, Kluin-Nelemans JC (2004): Gender and age-related differences in Burkitt lymphoma – epidemiological and clinical data from the Netherlands. *Eur J Cancer* 40:2781–2787.
- Boerma EG, Siebert R, Kluin PM, Baudis M (2009): Translocations involving 8q24 in Burkitt lymphoma and other malignant lymphomas: a historical review of cytogenetics in the light of today's knowledge. *Leukemia* 23:225–234.
- Bohers E, Mareschal S, Bouzeflen A, Marchand V, Ruminy P, Maingonnat C, et al. (2014): Targetable activating mutations are very frequent in GCB and ABC diffuse large B-cell lymphoma. *Genes Chromosomes Cancer* 53:144–153.
- Booman M, Douwes J, Glas AM, Riemersma SA, Jordanova ES, Kok K, et al. (2006): Mechanisms and effects of loss of human leukocyte antigen class II expression in immuneprivileged site-associated B-cell lymphoma. *Clin Cancer Res* 12:2698–2705.
- Booman M, Szuhai K, Rosenwald A, Hartmann E, Kluin-Nelemans H, de Jong D, et al. (2008): Genomic aberrations and gene expression in primary diffuse large B-cell lymphomas of immune-privileged sites: the importance of apoptosis and immunomodulatory pathways. *J Pathol* 216:209–217.



Bosga-Bouwer AG, van Imhoff GW, Boonstra R, van der Veen A, Haralambieva E, van den Berg A, et al. (2003): Follicular lymphoma grade 3B includes 3 cytogenetically defined subgroups with primary t(14;18), 3q27, or other translocations: t(14;18) and 3q27 are mutually exclusive. *Blood* 101:1149–1154.

Bosga-Bouwer AG, van den Berg A, Haralambieva E, de Jong D, Boonstra R, Kluin P, van den Berg E, Poppema S (2006): Molecular, cytogenetic, and immunophenotypic characterization of follicular lymphoma grade 3B; a separate entity or part of the spectrum of diffuse large B-cell lymphoma or follicular lymphoma? *Hum Pathol* 37:528–533.

Bosga-Bouwer AG, Haralambieva E, Booman M, Boonstra R, van den Berg A, Schuurin E, et al. (2005): BCL6 alternative translocation breakpoint cluster region associated with follicular lymphoma grade 3B. *Genes Chromosomes Cancer* 44:301–304.

Bouamar H, Abbas S, Lin AP, Wang L, Jiang D, Holder KN, et al. (2013): A capture-sequencing strategy identifies IRF8, EBF1, and APRIL as novel IGH fusion partners in B-cell lymphoma. *Blood* 122:726–733.

Boulanger E, Agbalika F, Maarek O, Daniel MT, Grollet L, Molina JM, et al. (2001): A clinical, molecular and cytogenetic study of 12 cases of human herpesvirus 8 associated primary effusion lymphoma in HIV-infected patients. *Hematol J* 2:172–179.

Bouska A, McKeithan TW, Deffenbacher KE, Lachel C, Wright GW, Iqbal J, et al. (2014): Genome-wide copynumber analyses reveal genomic abnormalities involved in transformation of follicular lymphoma. *Blood* 123:1681–1690.

Boxer LM, Dang CV (2001): Translocations involving c-myc and c-myc function. *Oncogene* 20: 5595–5610.

Buchonnet G, Lenain P, Ruminy P, Lepretre S, Stamatoullas A, Parmentier F, et al. (2000): Characterisation of BCL2-JH rearrangements

in follicular lymphoma: PCR detection of 3' BCL2 breakpoints and evidence of a new cluster. *Leukemia* 14:1563–1569.

Buchonnet G, Jardin F, Jean N, Bertrand P, Parmentier F, Tison S, et al. (2002): Distribution of BCL2 breakpoints in follicular lymphoma and correlation with clinical features: specific subtypes or same disease? *Leukemia* 16:1852–1856.

Busslinger M, Klix N, Pfeffer P, Graninger PG, Kozmik Z (1996): Deregulation of PAX-5 by translocation of the Emu enhancer of the IgH locus adjacent to two alternative PAX-5 promoters in a diffuse large-cell lymphoma. *Proc Natl Acad Sci USA* 93:6129–6134.

Butler MP, Iida S, Capello D, Rossi D, Rao PH, Nallasivam P, et al. (2002): Alternative translocation breakpoint cluster region 5' to BCL-6 in B-cell non-Hodgkin's lymphoma. *Cancer Res* 62:4089–4094.

Callanan MB, LeBaccon P, Mossuz P, Duley S, Bastard C, Hamoudi R, et al. (2000): The IgG Fc receptor, Fc gamma RIIB, is a target for deregulation by chromosomal translocation in malignant lymphoma. *Proc Natl Acad Sci USA* 97:309–314

Campo E (2012): New pathogenic mechanisms in Burkitt lymphoma. *Nat Genet* 44:1288–1289.

Campo E (2013): Whole genome profiling and other high throughput technologies in lymphoid neoplasms: current contributions and future hopes. *Mod Pathol* 26: S97–S110

Carbone A (2003): Emerging pathways in the development of AIDS-related lymphomas. *Lancet Oncol* 4: 22–29.

Chambwe N, Kormaksson M, Geng H, De S, Michor F, Johnson NA, et al. (2014): Variability in DNA methylation defines novel epigenetic subgroups of DLBCL associated with different clinical outcomes. *Blood* 123:1699–1708.

Chen W, Iida S, Louie DC, Dalla-Favera R, Chaganti RS (1998): Heterologous promoters fused to BCL6 by chromosomal transloca-

tions affecting band 3q27 cause its deregulated expression during B-cell differentiation. *Blood* 91:603–607.

Chen W, Palanisamy N, Schmidt H, Teruya-Feldstein J, Jhanwar SC, Zelenetz AD, et al. (2001a): Deregulation of FCGR2B expression by 1q21 rearrangements in follicular lymphomas. *Oncogene* 20:7686–7693.

Chen W, Itoyama T, Chaganti RS (2001b): Splicing factor SRP20 is a novel partner of BCL6 in a t(3;6)(q27;p21) translocation in transformed follicular lymphoma. *Genes Chromosomes Cancer* 32:281–284.

Chen YW, Hu XT, Liang AC, Au WY, So CC, Wong ML, et al. (2006): High BCL6 expression predicts better prognosis, independent of BCL6 translocation partner, or BCL6-deregulating mutations, in gastric lymphoma. *Blood* 108:2373–2383.

Cheung KJ, Delaney A, Ben-Neriah S, Schein J, Lee T, Shah SP, et al. (2010a): High resolution analysis of follicular lymphoma genomes reveals somatic recurrent sites of copy-neutral loss of heterozygosity and copy number alterations that target single genes. *Genes Chromosomes Cancer* 49:669–681.

Cheung KJ, Johnson NA, Affleck JG, Severnson T, Steidl C, Ben-Neriah S, et al. (2010b): Acquired TNFRSF14 mutations in follicular lymphoma are associated with worse prognosis. *Cancer Res* 70:9166–9174.

Cheung KJ, Rogic S, Ben-Neriah S, Boyle M, Connors JM, Gascoyne RD, et al. (2012): SNP analysis of minimally evolved t(14;18)(q32;q21)-positive follicular lymphomas reveals a common copy-neutral loss of heterozygosity pattern. *Cytogenet Genome Res* 136:38–43.

Chikatsu N, Kojima H, Suzukawa K, Shinagawa A, Nagasawa T, Ozawa H, et al. (2003): ALK+, CD30-, CD20- large Bcell lymphoma containing anaplastic lymphoma kinase (ALK) fused to clathrin heavy chain gene (CLTC). *Mod Pathol* 16:828–832.

Christie L, Kernohan N, Levison D, Sales M, Cunningham J, Gillespie K, et al. (2008): C-MYC translocation in t(14;18) positive follicular lymphoma at presentation: an adverse prognostic indicator? *Leuk Lymphoma* 49:470–476.

Coiffier B, Lepage E, Briere J, Herbrecht R, Tilly H, Bouabdallah R, et al. (2002): CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *N Engl J Med* 346:235–242.

Cong P, Raffeld M, Teruya-Feldstein J, Sorbara L, Pittaluga S, Jaffe ES (2002): In situ localization of follicular lymphoma: description and analysis by laser capture microdissection. *Blood* 99:3376–3382.

Corcoran MM, Mould SJ, Orchard JA, Ibbotson RE, Chapman RM, Boright AP, et al. (1999): Dysregulation of cyclin dependent kinase 6 expression in splenic marginal zone lymphoma through chromosome 7q translocations. *Oncogene* 18:6271–6277.

Cui J, Liu Q, Cheng Y, Chen S, Sun Q (2014): An intravascular large B-cell lymphoma with a t(3;14)(q27;q32) translocation. *J Clin Pathol* 67:279–281.

Dallery E, Galiege-Zouitina S, Collyn-d'Hooghe M, Quief S, Denis C, Hildebrand MP, et al. (1995): TTF, a gene encoding a novel small G protein, fuses to the lymphoma-associated LAZ3 gene by t(3;4) chromosomal translocation. *Oncogene* 10:2171–2178.

Dave SS, Wright G, Tan B, Rosenwald A, Gascoyne RD, Chan WC, et al. (2004): Prediction of survival in follicular lymphoma based on molecular features of tumor-infiltrating immune cells. *N Engl J Med* 351:2159–2169.

Dave SS, Fu K, Wright GW, Lam LT, Kluin P, Boerma EJ, et al. (2006): Molecular diagnosis of Burkitt's lymphoma. *N Engl J Med* 354:2431–2442.

De Oliveira FM, Rodrigues-Alves AP, Lucena-Araújo AR, de Paula Silva F, da Silva FB, Fal-

- cão RP (2014): Mantle cell lymphoma harboring Burkitt's-like translocations presents differential expression of aurora kinase genes compared with other 8q abnormalities. *Med Oncol* 31:931.
- Dewald GW, Brockman SR, Paternoster SF (2004): Molecular cytogenetic studies for hematological malignancies. *Cancer Treat Res* 121: 69–112.
- Dewald GW, Therneau T, Larson D, Lee YK, Fink S, Smoley S, et al. (2005): Relationship of patient survival and chromosome anomalies detected in metaphase and/or interphase cells at diagnosis of myeloma. *Blood* 106:3553–3558.
- Deweindt C, Kerckaert JP, Tilly H, Quief S, Nguyen VC, Bastard C (1993): Cloning of a breakpoint cluster region at band 3q27 involved in human non-Hodgkin's lymphoma. *Genes Chromosomes Cancer* 8:149–154.
- Dierlamm J, Baens M, Wlodarska I, Stefanova-Ouzounova M, Hernandez JM, Hossfeld DK, et al. (1999): The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood* 93:3601–3609.
- Dürig J, Bug S, Klein-Hitpass L, Boes T, Jöns T, Martin-Subero JI, et al. (2007): Combined single nucleotide polymorphism-based genomic mapping and global gene expression profiling identifies novel chromosomal imbalances, mechanisms and candidate genes important in the pathogenesis of T-cell prolymphocytic leukemia with inv(14)(q11q32). *Leukemia* 21:2153–2163.
- Dyer MJ (2003): The pathogenetic role of oncogenes deregulated by chromosomal translocation in B-cell malignancies. *Int J Hematol* 77:315–320.
- Dyer MJ (2013): The detection of chromosomal translocations involving the immunoglobulin loci in B-cell malignancies. *Methods Mol Biol* 971:123–133.
- Dyer MJ, Lillington DM, Bastard C, Tilly H, Lens D, Heward JM, et al. (1996): Concurrent activation of MYC and BCL2 in B cell non-Hodgkin lymphoma cell lines by translocation of both oncogenes to the same immunoglobulin heavy chain locus. *Leukemia* 10:1198–1208.
- Dyomin VG, Rao PH, Dalla-Favera R, Chaganti RS (1997): BCL8, a novel gene involved in translocations affecting band 15q11–13 in diffuse large-cell lymphoma. *Proc Natl Acad Sci USA* 94:5728–5732.
- Dyomin VG, Palanisamy N, Lloyd KO, Dyomina K, Jhanwar SC, Houldsworth J, et al. (2000): MUC1 is activated in a B-cell lymphoma by the t(1;14)(q21;q32) translocation and is rearranged and amplified in B-cell lymphoma subsets. *Blood* 95:2666–2671.
- Eide MB, Liestøl K, Lingjaerde OC, Hystad ME, Kresse SH, Meza-Zepeda L, et al. (2010): Genomic alterations reveal potential for higher grade transformation in follicular lymphoma and confirm parallel evolution of tumor cell clones. *Blood* 116:1489–1497.
- Einerson RR, Law ME, Blair HE, Kurtin PJ, McClure RF, Ketterling RP, et al. (2006): Novel FISH probes designed to detect IGH-MYC and IGL-MYC rearrangements in B-cell lineage malignancy identify a new breakpoint cluster region designated BVR2. *Leukemia* 20:1790–1799.
- Fenton JA, Vaandrager JW, Aarts WM, Bende RJ, Heering K, van Dijk M, et al. (2002): Follicular lymphoma with a novel t(14;18) breakpoint involving the immunoglobulin heavy chain switch mu region indicates an origin from germinal center B cells. *Blood* 99:716–718.
- Fenton JA, Pratt G, Rothwell DG, Rawstron AC, Morgan GJ (2004): Translocation t(11;14) in multiple myeloma: analysis of translocation breakpoints on der(11) and der(14) chromosomes suggests complex molecular mechanisms of recombination. *Genes Chromosomes Cancer* 39:151–155.
- Franke S, Wlodarska I, Maes B, Vandenberghe P, Achten R, Hagemeijer A, et al. (2002): Comparative genomic hybridization pattern distinguishes T-cell/histiocytic B-cell lymphoma from nodular lymphocyte predominance Hodgkin's lymphoma. *Am J Pathol* 161:1861–1867.
- Fu K, Weisenburger DD, Greiner TC, Dave S, Wright G, Rosenwald A, et al. (2005): Cyclin D1-negative mantle cell lymphoma: a clinicopathologic study based on gene expression profiling. *Blood* 106: 4315–4321.
- Fukuhara S, Rowley JD, Variakojis D, Golomb HM (1979): Chromosome abnormalities in poorly differentiated lymphocytic lymphoma. *Cancer Res* 39:3119–3128.
- Galiègue-Zouitina S, Quief S, Hildebrand MP, Denis C, Lecocq G, Collyn-d'Hooghe M, et al. (1996): The B cell transcriptional coactivator BOB1/OBF1 gene fuses to the LAZ3/BCL6 gene by t(3;11)(q27;q23.1) chromosomal translocation in a B cell leukemia line (Karpas 231). *Leukemia* 10:579–587.
- Gallieue-Zouitina S, Quief S, Hildebrand MP, Denis C, Detournignies L, Laï JL, et al. (1999): Nonrandom fusion of L-plastin(LCP1) and LAZ3(BCL6) gene by t(3;13) (q27;q14) chromosome translocation in two cases of Bcell non-Hodgkin lymphoma. *Genes Chromosomes Cancer* 26:97–105.
- Gascoyne RD, Lamant L, Martin-Subero JI, Lestou VS, Harris NL, Muller-Hermelink HK, et al. (2003): ALKpositive diffuse large B-cell lymphoma is associated with clathrin-ALK rearrangements: report of 6 cases. *Blood* 102:2568–2573.
- Gesk S, Gascoyne RD, Schnitzer B, Bakshi N, Janssen D, Klapper W, et al. (2005): ALK-positive diffuse large B-cell lymphoma with ALK-clathrin fusion belongs to the spectrum of pediatric lymphomas. *Leukemia* 19:1839–1840.
- Gesk S, Klapper W, Martin-Subero JI, Nagel I, Harder L, Fu K, et al. (2006): A chromosomal translocation in cyclin D1-negative/cyclin D2-positive mantle cell lymphoma fuses the CCND2 gene to the IGH locus. *Blood* 108:1109–1110.
- Giefing M, Arnemann J, Martin-Subero JI, Nieländer I, Bug S, Hartmann S, et al. (2008): Identification of candidate tumour suppressor gene loci for Hodgkin and Reed-Sternberg cells by characterisation of homozygous deletions in classical Hodgkin lymphoma cell lines. *Br J Haematol* 142:916–924.
- Gilles F, Goy A, Remache Y, Shue P, Zelenetz AD (2000): MUC1 dysregulation as the consequence of a t(1;14)(q21;q32) translocation in an extranodal lymphoma. *Blood* 95:2930–2936.
- Green MR, Gentles AJ, Nair RV, Irish JM, Kihira S, Liu CL, et al. (2013): Hierarchy in somatic mutations arising during genomic evolution and progression of follicular lymphoma. *Blood* 121:1604–1611.
- Green MR, Vicente-Dueñas C, Romero-Camarero I, Long Lui C, Dai B, Gonzalez-Herrero I, et al. (2014): Transient expression of BCL6 is sufficient for oncogenic function and induction of mature B-cell lymphoma. *Nat Commun* 5:3904.
- Gu K, Fu K, Jain S, Liu Z, Iqbal J, Li M, et al. (2009): t(14;18)-negative follicular lymphomas are associated with a high frequency of BCL6 rearrangement at the alternative breakpoint region. *Mod Pathol* 22:1251–1257.
- Gulia A, Saggini A, Wiesner T, Fink-Puches R, Argenyi Z, Ferrara G, et al. (2011): Clinicopathologic features of early lesions of primary cutaneous follicle center lymphoma, diffuse type: implications for early diagnosis and treatment. *J Am Acad Dermatol* 65:991–1000.
- Gutierrez-García G, Cardesa-Salzman T, Climent F, Gonzalez-Barca E, Mercadal S, Mate JL, et al. (2011): Gene-expression profiling and not immunophenotypic algorithms predict prognosis in patients with diffuse large B-cell lymphoma treated with immunochemotherapy. *Blood* 117:4836–4843.

Hallermann C, Kaune KM, Gesk S, Martin-Subero JI, Gunawan B, Griesinger F, et al. (2004a): Molecular cytogenetic analysis of chromosomal breakpoints in the IGH, MYC, BCL6, and MALT1 gene loci in primary cutaneous B-cell lymphomas. *J Invest Dermatol* 123:213–219.

Hallermann C, Kaune KM, Siebert R, Vermeer MH, Tensen CP, Willemze R, et al. (2004b): Chromosomal aberration patterns differ in subtypes of primary cutaneous B cell lymphomas. *J Invest Dermatol* 122:1495–1502.

Harris NL, Jaffe ES, Stein H, Banks PM, Chan JK, Cleary ML, et al. (1994): A revised European–American classification of lymphoid neoplasms: a proposal from the International Lymphoma Study Group. *Blood* 84:1361–1392.

Harris NL, Jaffe ES, Diebold J, Flandrin G, Muller-Hermelink HK, Vardiman J, et al. (1999): World Health Organization classification of neoplastic diseases of the hematopoietic and lymphoid tissues: report of the Clinical Advisory Committee Meeting, Airlie House, Virginia, November 1997. *J Clin Oncol* 17:3835–3849.

Harris NL, Jaffe ES, Diebold J, Flandrin G, Muller-Hermelink HK, Vardiman J, et al. (2000): The World Health Organization classification of hematological malignancies report of the Clinical Advisory Committee Meeting, Airlie House, Virginia, November 1997. *Mod Pathol* 13:193–207.

Harris NL, Swerdlow S, Jaffe ES, Ott G, Nathwani BN, de Jong D, et al. (2008): Follicular lymphoma. In: Swerdlow S, Campo E, Harris N, et al, eds. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues* IARC:220–226.

He L, Thomson JM, Hemann MT, Hernandez-Monge E, Mu D, Goodson S, et al. (2005): A microRNA polycistron as a potential human oncogene. *Nature* 435:828–833.

Herens C, Lambert F, Quintanilla-Martinez L, Bisig B, Deusings C, deLeval L (2008): Cyclin D1-negative mantle cell lymphoma with cryptic t(12;14)(p13;q32) and cyclin D2 overexpression. *Blood* 111:1745–1746.

Heyning FH, Jansen PM, Hogendoorn PC, Szuhai K (2010): Array-based comparative genomic hybridisation analysis reveals recurrent chromosomal alterations in primary diffuse large B cell lymphoma of bone. *J Clin Pathol* 63:1095–1100.

Hillion J, Mecucci C, Aventin A, Leroux D, Wlodarska I, Van Den Berghe H, et al. (1991): A variant translocation t(2;18) in follicular lymphoma involves the 5' end of bcl-2 and Igk light chain gene. *Oncogene* 6:169–172.

Höglund M, Sehn L, Connors JM, Gascoyne RD, Siebert R, Säll T, et al. (2004): Identification of cytogenetic subgroups and karyotypic pathways of clonal evolution in follicular lymphomas. *Genes Chromosomes Cancer* 39:195–204.

Honma K, Tsuzuki S, Nakagawa M, Karnan S, Aizawa Y, Kim WS, et al. (2008): TNFAIP3 is the target gene of chromosome band 6q23.3–q24.1 loss in ocular adnexal marginal zone B cell lymphoma. *Genes Chromosomes Cancer* 47:1–7.

Horn H, Schmelter C, Leich E, Salaverria I, Katzenberger T, Ott MM, et al. (2011): Follicular lymphoma grade 3B is a distinct neoplasm according to cytogenetic and immunohistochemical profiles. *Haematologica* 96:1327–1334.

Horsman D, Gascoyne R, Klasa R, Coupland R (1992): t(11;18)(q21;q21.1): a recurring translocation in lymphomas of mucosa-associated lymphoid tissue (MALT)? *Genes Chromosomes Cancer* 4:183–187.

Horsman DE, Connors JM, Pantzar T, Gascoyne RD (2001): Analysis of secondary chromosomal alterations in 165 cases of follicular lymphoma with t(14;18). *Genes Chromosomes Cancer* 30:375–382.

Horsman DE, Okamoto I, Ludkovski O, Le N, Harder L, Gesk S, et al. (2003): Follicular lymphoma lacking the t(14;18)(q32;q21): identification of two disease subtypes. *Br J Haematol* 120: 424–433.

Hosokawa Y, Maeda Y, Ichinohasama R, Miura I, Taniwaki M, Seto M (2000): The Ikaros gene,

a central regulator of lymphoid differentiation, fuses to the BCL6 gene as result of t(3;7)(q27;p12) translocation in a patient with diffuse large B-cell lymphoma. *Blood* 95:2719–2721.

Hu XT, Chen YW, Liang AC, Au WY, Wong KY, Wan TS, et al. (2012): CD44 activation in mature B-cell malignancies by a novel recurrent IGH translocation. *Blood* 115:2458–2461

Hummel M, Bentink S, Berger H, Klapper W, Wessendorf S, Barth TF, et al. (2006): A biologic definition of Burkitt's lymphoma from transcriptional and genomic profiling. *N Engl J Med* 354:2419–2430.

Iida S, Rao PH, Nallasivam P, Hibshoosh H, Butler M, Louie DC, et al. (1996): The t(9;14)(p13;q32) chromosomal translocation associated with lymphoplasmacytoid lymphoma involves the PAX-5 gene. *Blood* 88:4110–4117.

Iqbal J, Sanger WG, Horsman DE, Rosenwald A, Pickering DL, Dave B, et al. (2004): BCL2 translocation defines a unique tumor subset within the germinal center B-cell-like diffuse large B-cell lymphoma. *Am J Pathol* 165: 159–166.

Iqbal J, Greiner TC, Patel K, Dave BJ, Smith L, Ji J, et al. (2007): Distinctive patterns of BCL6 molecular alterations and their functional consequences in different subgroups of diffuse large B-cell lymphoma. *Leukemia* 21: 2332–2343.

Iqbal J, Kucuk C, Deleeuw RJ, Srivastava G, Tam W, Geng H, et al. (2009): Genomic analyses reveal global functional alterations that promote tumor growth and novel tumor suppressor genes in natural killer-cell malignancies. *Leukemia* 23:1139–1151.

Jaffe ES, Harris NL, Stein H, Vardiman JW, editors (2001): *The World Health Organization Classification of Tumours: Pathology & Genetics. Tumours of Haematopoietic and Lymphoid Tissues*. Lyon: IACR Press.

Jardin F, Gaulard P, Buchonnet G, Contentin N, Leprêtre S, Lenain P, et al. (2002): Follicular lymphoma without t(14;18) and with BCL-6 re-

arrangement: a lymphoma subtype with distinct pathological, molecular and clinical characteristics. *Leukemia* 16:2309–2317.

Jardin F, Ruminy P, Bastard C, Tilly H (2007): The BCL6 protooncogene: a leading role during germinal center development and lymphomagenesis. *Pathol Biol (Paris)* 55:73–83

Jares P, Colomer D, Campo E (2007): Genetic and molecular pathogenesis of mantle cell lymphoma: perspectives for new targeted therapeutics. *Nat Rev Cancer* 7:750–762.

Johansson B, Mertens F, Mitelman F (1995): Cytogenetic evolution patterns in non-Hodgkin's lymphoma. *Blood* 86:3905–3914.

Johnson NA, Al-Tourah A, Brown CJ, Connors JM, Gascoyne RD, Horsman DE (2008): Prognostic significance of secondary cytogenetic alterations in follicular lymphomas. *Genes Chromosomes Cancer* 47:1038–1048.

Johnson NA, Savage KJ, Ludkovski O, Ben-Neriah S, Woods R, Steidl C, et al. (2009): Lymphomas with concurrent BCL2 and MYC translocations: the critical factors associated with survival. *Blood* 114:2273–2279.

Joos S, Falk MH, Lichter P, Haluska FG, Henglein B, Lenoir GM, et al. (1992a): Variable breakpoints in Burkitt lymphoma cells with chromosomal t(8;14) translocation separate c-myc and the IgH locus up to several hundred kb. *Hum Mol Genet* 1:625–632.

Joos S, Haluska FG, Falk MH, Henglein B, Hameister H, Croce CM, et al. (1992b): Mapping chromosomal breakpoints of Burkitt's t(8;14) translocations far upstream of c-myc. *Cancer Res* 52:6547–6552.

Kaneita Y, Yoshida S, Ishiguro N, Sawada U, Horie T, Mori S, et al. (2001): Detection of reciprocal fusion 5'-BCL6/partner-3' transcripts in lymphomas exhibiting reciprocal BCL6 translocations. *Br J Haematol* 113:803–806.

Kanungo A, Medeiros LJ, Abruzzo LV, Lin P (2006): Lymphoid neoplasms associated with



- concurrent t(14;18) and 8q24/c-MYC translocation generally have a poor prognosis. *Mod Pathol* 19:25–33.
- Karsan A, Gascoyne RD, Coupland RW, Shepherd JD, Phillips GL, Horsman DE (1993): Combination of t(14;18) and a Burkitt's type translocation in B-cell malignancies. *Leuk Lymphoma* 10:433–441.
- Katzenger T, Ott G, Klein T, Kalla J, Muller-Hermelink HK, Ott MM (2004): Cytogenetic alterations affecting BCL6 are predominantly found in follicular lymphomas grade 3B with a diffuse large B-cell component. *Am J Pathol* 165: 481–490.
- Katzenger T, Kienle D, Stilgenbauer S, Höller S, Schilling C, Mäder U, et al. (2008): Delineation of distinct tumour profiles in mantle cell lymphoma by detailed cytogenetic, interphase genetic and morphological analysis. *Br J Haematol* 142:538–550.
- Katzenger T, Kalla J, Leich E, Stocklein H, Hartmann E, Barnickel S, et al. (2009): A distinctive subtype of t(14;18) negative nodal follicular non-Hodgkin lymphoma characterized by a predominantly diffuse growth pattern and deletions in the chromosomal region 1p36. *Blood* 113:1052–1061.
- Kearney L, Horsley SW (2005): Molecular cytogenetics in haematological malignancy: current technology and future prospects. *Chromosoma* 114: 286–294.
- Kerckaert JP, Deweindt C, Tilly H, Quief S, Lecocq G, Bastard C (1993): LAZ3, a novel zinc-finger encoding gene, is disrupted by recurring chromosome 3q27 translocations in human lymphomas. *Nat Genet* 5:66–70.
- Kim BK, Surti U, Pandya A, Cohen J, Rabkin MS, Swerdlow SH (2005): Clinicopathologic, immunophenotypic, and molecular cytogenetic fluorescence in situ hybridization analysis of primary and secondary cutaneous follicular lymphomas. *Am J Surg Pathol* 29:69–82.
- Kimm LR, deLeeuw RJ, Savage KJ, Rosenwald A, Campo E, Delabie J, et al. (2007): Frequent occurrence of deletions in primary mediastinal B-cell lymphoma. *Genes Chromosomes Cancer* 46:1090–1097.
- Klapper W, Szczepanowski M, Burkhardt B, Berger H, Rosolowski M, Bentink S, et al. (2008a): Molecular profiling of pediatric mature B-cell lymphoma treated in populationbased prospective clinical trials. *Blood* 112:1374–1381.
- Klapper W, Stoecklein H, Zeynalova S, Ott G, Kosari F, Rosenwald A, et al. (2008b): Structural aberrations affecting the MYC locus indicate a poor prognosis independent of clinical risk factors in diffuse large B-cell lymphomas treated within randomized trials of the German High-Grade Non-Hodgkin's Lymphoma Study Group (DSHNHL). *Leukemia* 22:2226–2229.
- Klapproth K, Wirth T (2010): Advances in the understanding of MYC-induced lymphomagenesis. *Br J Haematol* 149:484–497.
- Kluin PM (2013): Origin and migration of follicular lymphoma cells. *Haematologica* 98:1331–1333.
- Kluin PM (2014): The missing link in early follicular lymphoma development. *Blood* 123:3371–3372.
- Kluin P, Schuurin E (2011): Molecular cytogenetics of lymphoma: where do we stand in 2010? *Histopathology* 58: 128–144.
- Kluin PM, Harris NL, Stein H, Leoncini L, Raphaël M, Campo E, et al. (2008): B-cell lymphoma, unclassifiable, with features intermediate between diffuse large B-cell lymphoma and Burkitt lymphoma. In: Swerdlow S, Campo E, Harris N, Jaffe ES, Pileri S, Stein H, et al., eds. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. IARC, Lyon:265–266.
- Kluin-Nelemans HC, Hoster E, Hermine O, Walewski J, Trneny M, Geisler CH, et al. (2012): Treatment of older patients with mantle-cell lymphoma. *N Engl J Med* 367:520–531.
- Kneba M, Eick S, Herbst H, Willigeroth S, Pott C, Bolz I, et al. (1991): Frequency and structure of t(14;18) major breakpoint regions in non-Hodgkin's lymphomas typed according to the Kiel classification: analysis by direct DNA sequencing. *Cancer Res* 51:3243–3250.
- Knezevich S, Ludkovski O, Salski C, Lestou V, Chhanabhai M, Lam W, et al. (2005): Concurrent translocation of BCL2 and MYC with a single immunoglobulin locus in high-grade B-cell lymphomas. *Leukemia* 19:659–663.
- Knight SJ, Yau C, Clifford R, Timbs AT, Sadighi Akha E, Dréau HM, et al. (2012): Quantification of subclonal distributions of recurrent genomic aberrations in paired pre-treatment and relapse samples from patients with B-cell chronic lymphocytic leukemia. *Leukemia* 26:1564–1575.
- Kobayashi S, Taki T, Chinen Y, Tsutsumi Y, Ohshiro M, Kobayashi T, et al. (2011): Identification of IGH $\kappa$ -BACH2 fusion transcripts resulting from cryptic chromosomal rearrangements of 14q32 with 6q15 in aggressive B-cell lymphoma/leukemia. *Genes Chromosomes Cancer* 50:207–216.
- Kreisel F, Kulkarni S, Kerns RT, Hassan A, Deshmukh H, Nagarajan R, et al. (2011): High resolution array comparative genomic hybridization identifies copy number alterations in diffuse large B-cell lymphoma that predict response to immunochemotherapy. *Cancer Genet* 204:129–137.
- Kroenlein H, Schwartz S, Reinhardt R, Rieder H, Molkentin M, Gökbüget N, et al. (2012): Molecular analysis of the t(2;8)/MYC-IGK translocation in high-grade lymphoma/leukemia by long-distance inverse PCR. *Genes Chromosomes Cancer* 51:290–299.
- Kuehl WM, Bergsagel PL (2002): Multiple myeloma: evolving genetic events and host interactions. *Nat Rev Cancer* 2:175–187.
- Küppers R (2005): Mechanisms of B-cell lymphoma pathogenesis. *Nat Rev Cancer* 5: 251–262.
- Küppers R, Dalla-Favera R (2001): Mechanisms of chromosomal translocations in B cell lymphomas. *Oncogene* 20:5580–5594.
- Küppers R, Klein U, Hansmann ML, Rajewsky K (1999): Cellular origin of human B-cell lymphomas. *N Engl J Med* 341:1520–1529.
- Kurata M, Maesako Y, Ueda C, Nishikori M, Akasaka T, Uchiyama T, et al. (2002): Characterization of t(3;6)(q27;p21) breakpoints in B-cell non-Hodgkin's lymphoma and construction of the histone H4/BCL6 fusion gene, leading to altered expression of BCL6. *Cancer Res* 62:6224–6230.
- Kwiecinska A, Ichimura K, Berglund M, Dinets A, Sulaiman L, Collins VP, et al. (2014): Amplification of 2p as a genomic marker for transformation in lymphoma. *Genes Chromosomes Cancer* 53:750–768.
- Landau DA, Carter SL, Stojanov P, McKenna A, Stevenson K, Lawrence MS, et al. (2013): Evolution and impact of subclonal mutations in chronic lymphocytic leukemia. *Cell* 152:714–726.
- Lecointe N, Meerabux J, Ebihara M, Hill A, Young BD (1999): Molecular analysis of an unstable genomic region at chromosome band 11q23 reveals a disruption of the gene encoding the alpha2 subunit of platelet-activating factor acetylhydrolase (Pafah1a2) in human lymphoma. *Oncogene* 18:2852–2859.
- Leich E, Salaverria I, Bea S, Zettl A, Wright G, Moreno V, et al. (2009): Follicular lymphomas with and without translocation t(14;18) differ in gene expression profiles and genetic alterations. *Blood* 114:826–834.
- Lennert K, Stein H, Kaiserling E (1975): Cytological and functional criteria for the classification of malignant lymphomata. *Br J Cancer* 2:29–43.
- Lenz G, Nagel I, Siebert R, Roschke AV, Sanger W, Wright GW, et al. (2007): Aberrant immunoglobulin class switch recombination and switch translocations in activated B cell-like diffuse large B cell lymphoma. *J Exp Med* 204:633–643.

- Lenz G, Wright GW, Emre NC, Kohlhammer H, Dave SS, Davis RE, et al. (2008): Molecular subtypes of diffuse large B-cell lymphoma arise by distinct genetic pathways. *Proc Natl Acad Sci USA* 105:13520–13525.
- Leoncini L, Raphael M, Stein H, Harris NL, Jaffe ES, Kluin PM (2008): Burkitt lymphoma. In: Swerdlow S, Campo E, Harris N, Jaffe ES, Pileri S, Stein H, et al., eds. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. IARC, Lyon:262–264.
- Leroux D, Hillion J, Monteil M, Le Marc'hadour F, Jacob MC, Sotto JJ, et al. (1991): t(18;22) (q21;q11) with rearrangement of BCL2 as a possible secondary change in a lymphocytic lymphoma. *Genes Chromosomes Cancer* 3:205–209.
- Letai AG (2008): Diagnosing and exploiting cancer's addiction to blocks in apoptosis. *Nat Rev Cancer* 8:121–132.
- Leucci E, Cocco M, Onnis A, De Falco G, van Cleef P, Bellan C, et al. (2008): MYC translocation-negative Burkitt lymphoma cases: an alternative pathogenic mechanism involving miRNA deregulation. *J Pathol* 216:440–450.
- Limpens J, de Jong D, van Krieken JH, Price CG, Young BD, van Ommen GJ, et al. (1991): Bcl-2/JH rearrangements in benign lymphoid tissues with follicular hyperplasia. *Oncogene* 6:2271–2276.
- Limpens J, Stad R, Vos C, de Vlaam C, de Jong D, van Ommen GJ, et al. (1995): Lymphoma-associated translocation t(14;18) in blood B cells of normal individuals. *Blood* 85:2528–2536.
- Lin P, Jetly R, Lennon PA, Abruzzo LV, Prajapati S, Medeiros LJ (2008): Translocation (18;22) (q21;q11) in B-cell lymphomas: a report of 4 cases and review of literature. *Hum Pathol* 39:1664–1672.
- Liu Q, Salaverria I, Pittaluga S, Jegalian AG, Xi L, Siebert R, et al. (2013): Follicular lymphomas in children and young adults. *Am J Surg Pathol* 37:333–343.
- Loeffler M, Kreuz M, Haake A, Hasenclever D, Trautmann H, Arnold C (2015): Genomic and epigenomic co-evolution in follicular lymphomas. *Leukemia* 29:456–463.
- Lohr JG, Stojanov P, Lawrence MS, Auclair D, Chapuy B, Sougnez C, et al. (2012): Discovery and prioritization of somatic mutations in diffuse large B-cell lymphoma (DLBCL) by whole-exome sequencing. *Proc Natl Acad Sci USA* 109:3879–3884.
- Louissaint A Jr, Ackerman AM, Dias-Santagata D, Ferry JA, Hochberg EP, Huang MS, et al. (2012): Pediatric-type nodal follicular lymphoma: an indolent clonal proliferation in children and adults with high proliferation index and no BCL2 rearrangement. *Blood* 120:2395–2404.
- Love C, Sun Z, Jima D, Li G, Zhang J, Miles R, et al. (2012): The genetic landscape of mutations in Burkitt lymphoma. *Nat Genet* 44:1321–1325.
- Luan SL, Boulanger E, Ye H, Chanudet E, Johnson N, Hamoudi RA, et al. (2010): Primary effusion lymphoma: genomic profiling revealed amplification of SELPLG and CORO1C encoding for proteins important for cell migration. *J Pathol* 222:166–179.
- Maes B, Vanhentenrijk V, Wlodarska I, Cools J, Peeters B, Marynen P, et al. (2001): The NPM-ALK and the ATIC-ALK fusion genes can be detected in non-neoplastic cells. *Am J Pathol* 158:2185–2193.
- Majid A, Tsoulakis O, Walewska R, Gesk S, Siebert R, Kennedy DB, et al. (2007): BCL2 expression in chronic lymphocytic leukemia: lack of association with the BCL2 938A>C promoter single nucleotide polymorphism. *Blood* 111:874–877.
- Mamessier E, Song JY, Eberle FC, Pack S, Drevet C, Chetaille B, et al. (2014): Early lesions of follicular lymphoma: a genetic perspective. *Haematologica* 99:481–488.
- Mandal AK, Savvidou L, Slater RM, Cockett W, Wiggins J, Missouri CG (2007): Angiotropic lymphoma: associated chromosomal abnormalities. *Eur J Intern Med* 18:432–434.
- Manolov G, Manolova Y (1972): Marker band in one chromosome 14 from Burkitt lymphomas. *Nature* 237:33–34.
- Manolova Y, Manolov G, Kieler J, Levan A, Klein G (1979): Genesis of the 14q + marker in Burkitt's lymphoma. *Hereditas* 90:5–10.
- Martin LD, Belch AR, Pilarski LM (2010): Promiscuity of translocation partners in multiple myeloma. *J Cell Biochem* 109:1085–1094.
- Martin-Guerrero I, Salaverria I, Burkhardt B, Szczepanowski M, Baudis M, Bens S, et al. (2013): Recurrent loss of heterozygosity in 1p36 associated with TNFRSF14 mutations in IRF4 translocation negative pediatric follicular lymphoma. *Haematologica* 98:1237–1241.
- Martin-Subero JI, Gesk S, Harder L, Grote W, Siebert R (2003a): Interphase cytogenetics of hematological neoplasms under the perspective of the novel WHO classification. *Anticancer Res* 23:1139–1148.
- Martin-Subero JI, Knippschild U, Harder L, Barth TF, Riemke J, Grohmann S, et al. (2003b): Segmental chromosomal aberrations and centrosome amplifications: pathogenetic mechanisms in Hodgkin and Reed–Sternberg cells of classical Hodgkin's lymphoma? *Leukemia* 17:2214–2219.
- Martin-Subero JI, Otero MD, Hernandez R, Cigudosa JC, Agirre X, Saez B, et al. (2005): Amplification of IGH/MYC fusion in clinically aggressive IGH/BCL2-positive germinal center B-cell lymphomas. *Genes Chromosomes Cancer* 43:414–423.
- Masque-Soler N, Szczepanowski M, Kohler CW, Spang R, Klapper W (2013): Molecular classification of mature aggressive B-cell lymphoma using digital multiplexed gene expression on formalin-fixed paraffin-embedded biopsy specimens. *Blood* 122:1985–1986.
- May PC, Foot N, Dunn R, Geoghegan H, Neat MJ (2010): Detection of cryptic and variant IGH-MYC rearrangements in high-grade non-Hodgkin's lymphoma by fluorescence in situ hybridization: implications for cytogenetic testing. *Cancer Genet Cytogenet* 198:71–75.
- Micci F, Panagopoulos I, Tjønnfjord GE, Kolstad A, Delabie J, Beiske K, et al. (2007): Molecular cytogenetic characterization of t(14;19) (q32;p13), a new recurrent translocation in B cell malignancies. *Virchows Arch* 450:559–565.
- Miki T, Kawamata N, Hirose S, Aoki N (1994a): Gene involved in the 3q27 translocation associated with B-cell lymphoma, BCL5, encodes a Kruppel-like zinc-finger protein. *Blood* 83:26–32.
- Miki T, Kawamata N, Arai A, Ohashi K, Nakamura Y, Kato A, et al. (1994b): Molecular cloning of the breakpoint for 3q27 translocation in B-cell lymphomas and leukemias. *Blood* 83:217–222.
- Mirza I, Macpherson N, Paproski S, Gascoyne RD, Yang B, Finn WG, et al. (2002): Primary cutaneous follicular lymphoma: an assessment of clinical, immunophenotypic, and molecular features. *J Clin Oncol* 20:647–655.
- Mitelman F, Johansson B, Mertens F (Eds.) (2014): Mitelman Database of Chromosome Aberrations and Gene Fusions in Cancer. Available at <http://cgap.nci.nih.gov/Chromosomes/Mitelman>.
- Miyoshi I, Hiraki S, Kimura I, Miyamoto K, Sato J (1979): 2/8 translocation in a Japanese Burkitt's lymphoma. *Experientia* 35:742–743.
- Montesinos-Rongen M, Akasaka T, Zuhlke-Jenisch R, Schaller C, Van Roost D, et al. (2003): Molecular characterization of BCL6 breakpoints in primary diffuse large B-cell lymphoma of the central nervous system identifies GAPD as novel translocation partner. *Brain Pathol* 13:534–538.



- Montoto S, Fitzgibbon J (2011): Transformation of indolent B-cell lymphomas. *J Clin Oncol* 29:1827–1834.
- Montoto S, Davies AJ, Matthews J, Calaminici M, Norton AJ, Amess J, et al. (2007): Risk and clinical implications of transformation of follicular lymphoma to diffuse large B-cell lymphoma. *J Clin Oncol* 25:2426–2433.
- Morin RD, Mendez-Lago M, Mungall AJ, Goya R, Mungall KL, Corbett RD, et al. (2011): Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma. *Nature* 476:298–303.
- Morin RD, Mungall K, Pleasance E, Mungall AJ, Goya R, Huff RD, et al. (2013): Mutational and structural analysis of diffuse large B-cell lymphoma using whole-genome sequencing. *Blood* 122:1256–1265.
- Morris SW, Kirstein MN, Valentine MB, Dittmer KG, Shapiro DN, Saltman DL, et al. (1994): Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science* 263:1281–1284.
- Motokura T, Bloom T, Kim HG, Jüppner H, Ruderman JV, Kronenberg HM, et al. (1991): A novel cyclin encoded by a bcl1-linked candidate oncogene. *Nature* 350: 512–515.
- Mullaney BP, Ng VL, Herndier BG, McGrath MS, Pallavicini MG (2000): Comparative genomic analyses of primary effusion lymphoma. *Arch Pathol Lab Med* 124: 824–826.
- Mwenifumbo JC, Marra MA (2013): Cancer genomesequencing study design. *Nat Rev Genet* 14:321–332.
- Nadal N, Chapiro E, Flandrin-Gresta P, Thouvenin S, Vasselon C, Beldjord K, et al. (2012): LHX2 deregulation by juxtaposition with the IGH locus in a pediatric case of chronic myeloid leukemia in B-cell lymphoid blast crisis. *Leuk Res* 36:e195–e198.
- Nagel I, Bug S, Tönnies H, Ammerpohl O, Richter J, Vater I, et al. (2009a): Biallelic inactivation of TRAF3 in a subset of B-cell lymphomas with interstitial del(14)(q24.1q32.33) *Leukemia* 23:2153–2155.
- Nagel I, Akasaka T, Klapper W, Gesk S, Böttcher S, Ritgen M, et al. (2009b): Identification of the gene encoding cyclin E1 (CCNE1) as a novel IGH translocation partner in t(14;19)(q32;q12) in diffuse large B-cell lymphoma. *Haematologica* 94:1020–1023.
- Nakamura Y, Takahashi N, Kakegawa E, Yoshida K, Ito Y, Kayano H, et al. (2008): The GAS5 (growth arrest-specific transcript 5) gene fuses to BCL6 as a result of t(1;3)(q25;q27) in a patient with B-cell lymphoma. *Cancer Genet Cytogenet* 182:144–149.
- Navarro A, Clot G, Royo C, Jares P, Hadzidimitriou A, Agathangelidis A, et al. (2012): Molecular subsets of mantle cell lymphoma defined by the IGHV mutational status and SOX11 expression have distinct biologic and clinical features. *Cancer Res* 72:5307–5316.
- Nelson M, Perkins SL, Dave BJ, Coccia PF, Bridge JA, Lyden ER, et al. (2010): An increased frequency of 13q deletions detected by fluorescence in situ hybridization and its impact on survival in children and adolescents with Burkitt lymphoma: results from the Children's Oncology Group study CCG-5961. *Br J Haematol* 148:600–610.
- Neri A, Chang CC, Lombardi L, Salina M, Corradini P, Maiolo AT, et al. (1991): B cell lymphoma-associated chromosomal translocation involves candidate oncogene *lyt-10*, homologous to NF-kappa B p50. *Cell* 67:1075–1087.
- Ngo VN, Young RM, Schmitz R, Jhavar S, Xiao W, Lim KH, et al. (2011): Oncogenetically active MYD88 mutations in human lymphoma. *Nature* 470:115–119.
- Nieländer I, Bug S, Richter J, Giefing M, Martin-Subero JL, Siebert R (2007): Combining array-based approaches for the identification of candidate tumor suppressor loci in mature lymphoid neoplasms. *APMIS* 115:1107–1134.
- Niitsu N, Okamoto M, Nakamura N, Nakamine H, Aoki S, Hirano M, et al. (2007): Prognostic impact of chromosomal alteration of 3q27 on nodal B-cell lymphoma: correlation with histology, immunophenotype, karyotype, and clinical outcome in 329 consecutive patients. *Leuk Res* 31:1191–1197.
- Nowakowski GS, Dewald GW, Hoyer JD, Pateroster SF, Stockero KJ, Fink SR, et al. (2005): Interphase fluorescence in situ hybridization with an IGH probe is important in the evaluation of patients with a clinical diagnosis of chronic lymphocytic leukaemia. *Br J Haematol* 130:36–42.
- O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT (2005): c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 435:839–843.
- O'Shea D, O'Riain C, Taylor C, Waters R, Carloti E, Macdougall F, et al. (2008): The presence of TP53 mutation at diagnosis of follicular lymphoma identifies a high-risk group of patients with shortened time to disease progression and poorer overall survival. *Blood* 112:3126–3129.
- Offit K, Parsa NZ, Gaidano G, Filippa DA, Louie D, Pan D, et al. (1993): 6q deletions define distinct clinico-pathologic subsets of non-Hodgkin's lymphoma. *Blood* 82:2157–2162.
- Ohno H (2004): Pathogenetic role of BCL6 translocation in B-cell non-Hodgkin's lymphoma. *Histol Histopathol* 19:637–650.
- Ohno H (2006): Pathogenetic and clinical implications of non-immunoglobulin; BCL6 translocations in B-cell non-Hodgkin's lymphoma. *J Clin Exp Hematop* 46:43–53.
- Ohno H, Takimoto G, McKeithan TW (1990): The candidate proto-oncogene bcl-3 is related to genes implicated in cell lineage determination and cell cycle control. *Cell* 60:991–997.
- Okosun J, Bödder C, Wang J, Araf S, Yang CY, Pan C, et al. (2014): Integrated genomic analysis identifies recurrent mutations and evolution patterns driving the initiation and progression of follicular lymphoma. *Nat Genet* 46:176–181.
- Oschlies I, Salaverria I, Mahn F, Meinhardt A, Zimmermann M, Woessmann W, et al. (2010): Pediatric follicular lymphoma—a clinico-pathological study of a populationbased series of patients treated within the Non-Hodgkin's Lymphoma – Berlin-Frankfurt-Münster (NHL-BFM) multicenter trials. *Haematologica* 95:253–259.
- Ott G, Katzenberger T, Greiner A, Kalla J, Rosenwald A, Heinrich U, et al. (1997): The t(11;18)(q21;q21) chromosome translocation is a frequent and specific aberration in low-grade but not high-grade malignant non-Hodgkin's lymphomas of the mucosa-associated lymphoid tissue (MALT-) type. *Cancer Res* 57:3944–3948.
- Ott G, Katzenberger T, Lohr A, Kindelberger S, Rüdiger T, Wilhelm M, et al. (2002): Cytomorphologic, immunohistochemical, and cytogenetic profiles of follicular lymphoma: 2 types of follicular lymphoma grade 3. *Blood* 99:3806–3812.
- Ott G, Rosenwald A, Campo E (2013): Understanding MYC-driven aggressive B-cell lymphomas: pathogenesis and classification. *Blood* 122:3884–3891.
- Parker E, Macdonald JR, Wang C (2011): Molecular characterization of a t(2;7) translocation linking CDK6 to the IGH locus in CD5(–) monoclonal B-cell lymphocytosis. *Cancer Genet* 204:260–264.
- Parker EP, Siebert R, Oo TH, Schneider D, Hayette S, Wang C (2013): Sequencing of t(2;7) translocations reveals a consistent breakpoint linking CDK6 to the IGH locus in indolent B-cell neoplasia. *J Mol Diagn* 15:101–109.
- Pasqualucci L, Neumeister P, Goossens T, Nanjangud G, Chaganti RS, Küppers R, et al. (2001): Hypermutation of multiple proto-oncogenes in B-cell diffuse large-cell lymphomas. *Nature* 412:341–346.
- Pasqualucci L, Migliazza A, Basso K, Houldsworth J, Chaganti RS, Dalla-Favera R (2003):

Mutations of the BCL6 proto-oncogene disrupt its negative autoregulation in diffuse large B-cell lymphoma. *Blood* 101:2914–2923.

Pasqualucci L, Compagno M, Houldsworth J, Monti S, Grunn A, Nandula SV, et al. (2006): Inactivation of the PRDM1/BLIMP1 gene in diffuse large B cell lymphoma. *J Exp Med* 203: 311–317.

Pasqualucci L, Trifonov V, Fabbri G, Ma J, Rossi D, Chiarenza A, et al. (2011a): Analysis of the coding genome of diffuse large B-cell lymphoma. *Nat Genet* 43:830–837.

Pasqualucci L, Dominguez-Sola D, Chiarenza A, Fabbri G, Grunn A, Trifonov V, et al. (2011b): Inactivating mutations of acetyltransferase genes in B-cell lymphoma. *Nature* 471:189–195.

Pasqualucci L, Khiabani H, Fangazio M, Vassishtha M, Messina M, Holmes AB, et al. (2014): Genetics of follicular lymphoma transformation. *Cell Rep* 6:130–140.

Poirel HA, Cairo MS, Heerema NA, Swansbury J, Aupérin A, Launay E, et al. (2009): Specific cytogenetic abnormalities are associated with a significantly inferior outcome in children and adolescents with mature B-cell non-Hodgkin's lymphoma: results of the FAB/LMB 96 international study. *Leukemia* 23:323–331.

Pospisilova H, Baens M, Michaux L, Stul M, Van Hummelen P, Van Loo P, et al. (2007): Interstitial del(14)(q) involving IGH: a novel recurrent aberration in B-NHL. *Leukemia* 21:2079–2083.

Puente XS, Pinyol M, Quesada V, Conde L, Ordóñez GR, Villamor N, et al. (2011): Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature* 475:101–105.

Put N, Van Roosbroeck K, Konings P, Meeus P, Brusselmans C, Rack K, et al. (2012): Chronic lymphocytic leukemia and prolymphocytic leukemia with MYC translocations: a subgroup with an aggressive disease course. *Ann Hematol* 91:863–873.

Rashid R, Johnson RJ, Morris S, Dickinson H, Czyz J, O'Connor SJ, et al. (2006): Intravascular large B-cell lymphoma associated with a near-tetraploid karyotype, rearrangement of BCL6, and a t(11;14)(q13;q32). *Cancer Genet Cytogenet* 171:101–104.

Richter J, Schlesner M, Hoffmann S, Kreuz M, Leich E, Burkhardt B, et al. (2012): Recurrent mutations of the ID3 gene in Burkitt lymphoma identified by integrated genome, exome and transcriptome sequencing. *Nat Genet* 44:1316–1320.

Riemersma SA, Jordanova ES, Schop RF, Philippo K, Looijenga LH, Schuurin E, et al. (2000): Extensive genetic alterations of the HLA region, including homozygous deletions of HLA class II genes in B-cell lymphomas arising in immune-privileged sites. *Blood* 96:3569–3577.

Rosenberg CL, Wong E, Petty EM, Bale AE, Tsujimoto Y, Harris NL, et al. (1991): PRAD1, a candidate BCL1 oncogene: mapping and expression in centrocytic lymphoma. *Proc Natl Acad Sci USA* 88:9638–9642.

Rosenwald A, Wright G, Chan WC, Connors JM, Campo E, Fisher RI, et al. (2002): The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med* 346:1937–1947.

Rosnwald A, Wright G, Leroy K, Yu X, Gaulard P, Gascoyne RD, et al. (2003): Molecular diagnosis of primary mediastinal B cell lymphoma identifies a clinically favorable subgroup of diffuse large B cell lymphoma related to Hodgkin lymphoma. *J Exp Med* 198:851–862.

Ross CW, Ouillette PD, Saddler CM, Shedden KA, Malek SN (2007): Comprehensive analysis of copy number and allele status identifies multiple chromosome defects underlying follicular lymphoma pathogenesis. *Clin Cancer Res* 13:4777–4785.

Rossi D, Khiabani H, Spina V, Ciardullo C, Bruscaggini A, Famà R, et al. (2014): Clinical impact of small TP53 mutated subclones in chronic lymphocytic leukemia. *Blood* 123:2139–2147.

Roulland S, Lebailly P, Lecluse Y, Briand M, Potier D, Gauduchon P (2004): Characterization of the t(14;18) BCL2-IGH translocation in farmers occupationally exposed to pesticides. *Cancer Res* 64:2264–2269.

Roulland S, Lebailly P, Lecluse Y, Heutte N, Nadel B, Gauduchon P (2006a): Long-term clonal persistence and evolution of t(14;18)-bearing B cells in healthy individuals. *Leukemia* 20:158–162.

Roulland S, Navarro JM, Grenot P, Milili M, Agopian J, Montpelliér B, et al. (2006b): Follicular lymphoma-like B cells in healthy individuals: a novel intermediate step in early lymphomagenesis. *J Exp Med* 203:2425–2431.

Roulland S, Kelly RS, Morgado E, Sungalee S, Solal-Celigny P, Colombat P, Jouve N, et al. (2014): t(14;18) translocation: a predictive blood biomarker for follicular lymphoma development. *J Clin Oncol* 32:1347–1355.

Ruminy P, Jardin F, Picquenot JM, Gaulard P, Parmentier F, Buchonnet G, et al. (2006): Two patterns of chromosomal breakpoint locations on the immunoglobulin heavy-chain locus in B-cell lymphomas with t(3;14)(q27;q32): relevance to histology. *Oncogene* 25:4947–4954.

Salaverria I, Siebert R (2011): The gray zone between Burkitt's lymphoma and diffuse large B-cell lymphoma from a genetics perspective. *J Clin Oncol* 29:1835–1843.

Salaverria I, Zettl A, Beà S, Hartmann EM, Dave SS, Wright GW, et al. (2008): Chromosomal alterations detected by comparative genomic hybridization in subgroups of gene expression-defined Burkitt's lymphoma. *Haematologica* 93:1327–1334.

Salaverria I, Philipp C, Oschlies I, Kohler CW, Kreuz M, Szczepanowski M, et al. (2011): Translocations activating IRF4 identify a subtype of germinal center-derived B-cell lymphoma affecting predominantly children and young adults. *Blood* 118:139–147.

Salaverria I, Akasaka T, Gesk S, Szczepanowski M, Burkhardt B, Harder L, et al. (2012): The CBFA2T3/ACSF3 locus is recurrently involved in IGH chromosomal translocation t(14;16)(q32;q24) in pediatric B-cell lymphoma with germinal center phenotype. *Genes Chromosomes Cancer* 51:338–343.

Salaverria I, Martin-Guerrero I, Burkhardt B, Kreuz M, Zenz T, Oschlies I, et al. (2013): High resolution copy number analysis of IRF4 translocation-positive diffuse large B-cell and follicular lymphomas. *Genes Chromosomes Cancer* 52:150–155.

Salaverria I, Martin-Guerrero I, Wagener R, Kreuz M, Kohler CW, Richter J, et al. (2014): A recurrent 11q aberration pattern characterizes a subset of MYC-negative high-grade B-cell lymphomas resembling Burkitt lymphoma. *Blood* 123:1187–1198.

Sanchez-Izquierdo D, Buchonnet G, Siebert R, Gascoyne RD, Climent J, Karan L, et al. (2003): MALT1 is deregulated by both chromosomal translocation and amplification in B-cell non-Hodgkin lymphoma. *Blood* 101:4539–4546.

Santra M, Zhan F, Tian E, Barlogie B, Shaughnessy Jr (2003): A subset of multiple myeloma harboring the t(4;14)(p16;q32) translocation lacks FGFR3 expression but maintains an IGH/MMSET fusion transcript. *Blood* 101:2374–2376.

Satterwhite E, Sonoki T, Willis TG, Harder L, Nowak R, Arriola EL, et al. (2001): The BCL11 gene family: involvement of BCL11A in lymphoid malignancies. *Blood* 98:3413–3420.

Savage KJ, Monti S, Kutok JL, Cattoretto G, Neuberg D, et al. (2003): The molecular signature of mediastinal large B-cell lymphoma differs from that of other diffuse large B-cell lymphomas and shares features with classical Hodgkin lymphoma. *Blood* 102:3871–3879.

Savage KJ, Johnson NA, Ben-Neriah S, Connors JM, Sehn LH, Farinha P, et al. (2009): MYC gene rearrangements are associated with

a poor prognosis in diffuse large B-cell lymphoma patients treated with R-CHOP chemotherapy. *Blood* 114:3533–3537.

Schlegelberger B, Zwingers T, Harder L, Nowotny H, Siebert R, Vesely M, et al. (1999): Clinicopathogenetic significance of chromosomal abnormalities in patients with blastic peripheral B-cell lymphoma. *Blood* 94:3114–3120.

Schmidt J, Salaverria I, Haake A, Bonzheim I, Adam P, Montes-Moreno S, et al. (2014): Increasing genomic and epigenomic complexity in the clonal evolution from in situ to manifest t(14;18)-positive follicular lymphoma. *Leukemia* 28:1103–1112.

Schmitz R, Hansmann ML, Bohle V, Martin-Subero JI, Hartmann S, Mechttersheimer G, et al. (2009): TNFAIP3 (A20) is a tumor suppressor gene in Hodgkin lymphoma and primary mediastinal B cell lymphoma. *J Exp Med* 206:981–989.

Schmitz R, Young RM, Ceribelli M, Jhavar S, Xiao W, Zhang M, et al. (2012): Burkitt lymphoma pathogenesis and therapeutic targets from structural and functional genomics. *Nature* 490:116–120.

Schneider B, Nagel S, Kaufmann M, Winkelmann S, Bode J, Drexler HG, et al. (2008): t(3;7)(q27;q32) fuses BCL6 to a non-coding region at FRA7H near miR-29. *Leukemia* 22:1262–1266.

Schraders M, de Jong D, Kluin P, Groenen P, van Krieken H (2005): Lack of Bcl-2 expression in follicular lymphoma may be caused by mutations in the BCL2 gene or by absence of the t(14;18) translocation. *J Pathol* 205:329–335.

Schuh A, Becq J, Humphray S, Alexa A, Burns A, Clifford R, et al. (2012): Monitoring chronic lymphocytic leukemia progression by whole genome sequencing reveals heterogeneous clonal evolution patterns. *Blood* 120:4191–4196.

Schwaenen C, Wessendorf S, Kestler HA, Döhner H, Lichter P, Bentz M (2003): DNA microarray analysis in malignant lymphomas. *Ann Hematol* 82:323–332.

Schwaenen C, Viardot A, Berger H, Barth TF, Bentink S, Döhner H, et al. (2009): Microarray-based genomic profiling reveals novel genomic aberrations in follicular lymphoma which associate with patient survival and gene expression status. *Genes Chromosomes Cancer* 48:39–54.

Schwindt H, Akasaka T, Zuhlke-Jenisch R, Hans V, Schaller C, Klapper W, et al. (2006): Chromosomal translocations fusing the BCL6 gene to different partner loci are recurrent in primary central nervous system lymphoma and may be associated with aberrant somatic hypermutation or defective class switch recombination. *J Neuropathol Exp Neurol* 5:776–782.

Schwindt H, Vater I, Kreuz M, Montesinos-Rongen M, Brunn A, Richter J, et al. (2009): Chromosomal imbalances and partial uniparental disomies in primary central nervous system lymphoma. *Leukemia* 23:1875–1884.

Seifert M, Scholtysik R, Küppers R (2013): Origin and pathogenesis of B cell lymphomas. *Methods Mol Biol* 971:1–25.

Sen F, Lai R, Albitar M (2002): Chronic lymphocytic leukemia with t(14;18) and trisomy 12. *Arch Pathol Lab Med* 126:1543–1546.

Shimanuki M, Sonoki T, Hosoi H, Watanuki J, Murata S, Kawakami K, et al. (2013): Molecular cloning of IGL rearrangements using long-distance inverse PCR (LDI-PCR). *Eur J Haematol* 90:59–67.

Shustik J, Han G, Farinha P, Johnson NA, Ben Neriah S, Connors JM, et al. (2010): Correlations between BCL6 rearrangements and outcome in patients with diffuse large B-cell lymphoma treated with CHOP or R-CHOP. *Haematologica* 95:96–101.

Siebert R, Rosenwald A, Staudt LM, Morris SW (2001): Molecular features of B-cell lymphoma. *Curr Opin Oncol* 13:316–324.

Sonoki T, Harder L, Horsman DE, Karran L, Taniguchi I, Willis TG, et al. (2001): Cyclin D3 is a target gene of t(6;14)(p21.1;q32.3) of mature

B-cell malignancies. *Blood* 98:2837–2844.

Sonoki T, Willis TG, Oscier DG, Karran EL, Siebert R, Dyer MJ (2004): Rapid amplification of immunoglobulin heavy chain switch (IGHS) translocation breakpoints using long-distance inverse PCR. *Leukemia* 18:2026–2031.

Sonoki T, Tatetsu H, Nagasaki A, Hata H (2007): Molecular cloning of translocation breakpoint from der(8)t(3;8)(q27;q24) defines juxtaposition of downstream of C-MYC and upstream of BCL6. *Int J Hematol* 86:196–198.

Streubel B, Lamprecht A, Dierlamm J, Cerroni L, Stolte M, Ott G, et al. (2003): t(14;18)(q32;q21) involving IGH and MALT1 is a frequent chromosomal aberration in MALT lymphoma. *Blood* 101:2335–2339.

Streubel B, Chott A, Huber D, Exner M, Jäger U, Wagner O, et al. (2004): Lymphoma-specific genetic aberrations in microvascular endothelial cells in B-cell lymphomas. *N Engl J Med* 351:250–259.

Streubel B, Vinatzer U, Lamprecht A, Raderer M, Chott A (2005): t(3;14)(p14.1;q32) involving IGH and FOXP1 is a novel recurrent chromosomal aberration in MALT lymphoma. *Leukemia* 19:652–658.

Streubel B, Scheucher B, Valencak J, Huber D, Petzelbauer P, Trautinger F, et al. (2006a): Molecular cytogenetic evidence of t(14;18)(IGH;BCL2) in a substantial proportion of primary cutaneous follicle center lymphomas. *Am J Surg Pathol* 30:529–536.

Streubel B, Vinatzer U, Willheim M, Raderer M, Chott A (2006b): Novel t(5;9)(q33;q22) fuses ITK to SYK in unspecified peripheral T-cell lymphoma. *Leukemia* 20:313–318.

Suguro M, Yoshida N, Umino A, Kato H, Tagawa H, Nakagawa M, et al. (2014): Clonal heterogeneity of lymphoid malignancies correlates with poor prognosis. *Cancer Sci* 105:897–904.

Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et al. (Eds.) (2008): *WHO*

*Classification of Tumors of Haematopoietic and Lymphoid Tissues*. Lyon: IARC.

Tagawa H, Suguro M, Tsuzuki S, Matsuo K, Karan S, Ohshima K, et al. (2005): Comparison of genome profiles for identification of distinct subgroups of diffuse large B-cell lymphoma. *Blood* 106: 1770–1777 (Erratum in Blood, 2006, 107:3052).

Tam W, Gomez M, Chadburn A, Lee JW, Chan WC, Knowles DM (2006): Mutational analysis of PRDM1 indicates a tumor-suppressor role in diffuse large B-cell lymphomas. *Blood* 107:4090–4100.

Tanaka R, Satoh H, Moriyama M, Satoh K, Morishita Y, Yoshida S, et al. (2000): Intronic U50 small-nucleolar-RNA (snoRNA) host gene of no protein-coding potential is mapped at the chromosome breakpoint t(3;6)(q27;q15) of human B-cell lymphoma. *Genes Cells* 5:277–287.

Taub R, Kirsch I, Morton C, Lenoir G, Swan D, Tronick S, et al. (1982): Translocation of the c-myc gene into the immunoglobulin heavy chain locus in human Burkitt lymphoma and murine plasmacytoma cells. *Proc Natl Acad Sci USA* 79:7837–7841.

Tellier J, Menard C, Roulland S, Martin N, Monvoisin C, Chasson L, et al. (2014): Human t(14;18)positive germinal center B cells: a new step in follicular lymphoma pathogenesis? *Blood* 123:3462–3465.

The International Non-Hodgkin's Lymphoma Prognostic Factors Project (1993): A predictive model for aggressive non-Hodgkin's lymphoma. *N Engl J Med* 329:987–994.

Thomadaki H, Scorilas A (2006): BCL2 family of apoptosis-related genes: functions and clinical implications in cancer. *Crit Rev Clin Lab Sci* 43:1–67.

Tinguely M, Thies S, Frigerio S, Reineke T, Korol D, Zimmermann DR (2014): IRF8 is associated with germinal center B-cell-like type of diffuse large B-cell lymphoma and exceptionally



- translocation t(14;16)(q32.33;q24.1). *Leuk Lymphoma* 55:136–142.
- Tsujimoto Y, Finger LR, Yunis J, Nowell PC, Croce CM (1984): Cloning of the chromosome breakpoint of neoplastic B cells with the t(14;18) chromosome translocation. *Science* 226:1097–1099.
- Tsujimoto Y, Gorham J, Cossman J, Jaffe E, Croce CM (1985): The t(14;18) chromosome translocations involved in B-cell neoplasms result from mistakes in VDJ joining. *Science* 229:1390–1393.
- Tzankov A, Xu-Monette ZY, Gerhard M, Visco C, Dirnhofer S, Gisin N, et al. (2014): Rearrangements of MYC gene facilitate risk stratification in diffuse large B-cell lymphoma patients treated with rituximab-CHOP. *Mod Pathol* 27:958–971.
- Ueda C, Akasaka T, Ohno H (2002a): Non-immunoglobulin/BCL6 gene fusion in diffuse large B-cell lymphoma: prognostic implications. *Leuk Lymphoma* 43:1375–1381.
- Ueda C, Akasaka T, Kurata M, Maesako Y, Nishikori M, Ichinohasama R, et al. (2002b): The gene for interleukin-21 receptor is the partner of BCL6 in t(3;16)(q27;p11), which is recurrently observed in diffuse large B-cell lymphoma. *Oncogene* 21:368–376.
- Vaandrager JW, Schuurin E, Philippo K, Kluin PM (2000): V(D)J recombinase-mediated transposition of the BCL2 gene to the IGH locus in follicular lymphoma. *Blood* 96:1947–1952.
- Vaghefi P, Martin A, Prévot S, Charlotte F, Camilleri-Broët S, Barli E, et al. (2006): Genomic imbalances in AIDS-related lymphomas: relation with tumoral Epstein–Barr virus status. *AIDS* 20:2285–2291.
- Valera A, Coloma L, Martinez A, de Jong D, Balagué O, Matheu G, et al. (2013): ALK-positive large B-cell lymphomas express a terminal B-cell differentiation program and activated STAT3 but lack MYC rearrangements. *Mod Pathol* 26:1329–1337.
- Van Den Berghe H, Louwagie A, Broeckaert-Van Orshoven A, David G, Verwilghen R, Michaux JL, et al. (1979): Philadelphia chromosome in human multiple myeloma. *J Natl Cancer Inst* 63:11–16.
- Van Dongen JJ, Langerak AW, Brüggemann M, Evans PA, Hummel M, Lavender FL, et al. (2003): Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect lymphoproliferations: report of the BIOMED-2 Concerted Action BMH4-CT98-3936. *Leukemia* 17:2257–2317.
- Vaqué JP, Martínez N, Batlle-López A, Pérez C, Montes-Moreno S, Sanchez-Beato M, et al. (2014): B-cell lymphoma mutations: improving diagnostics and enabling targeted therapies. *Haematologica* 99:222–231.
- van der Velden VHJ, Hoogeveen PG, de Ridder D, Schnindler-van der Struijk M, van Zelm MC, Sanders M, et al. (2014): B-cell prolymphocytic leukemia: a specific subgroup of mantle cell lymphoma. *Blood* 124:412–419.
- Ventura RA, Martin-Subero JI, Jones M, McParland J, Gesk S, Mason DY, et al. (2006): FISH analysis for the detection of lymphoma-associated chromosomal abnormalities in routine paraffin-embedded tissue. *J Mol Diagn* 8:141–151.
- Vergier B, Belaud-Rotureau MA, Benassy MN, Beylot-Barry M, Dubus P, Delaunay M, et al. (2004): Neoplastic cells do not carry bcl2-JH rearrangements detected in a subset of primary cutaneous follicle center B-cell lymphomas. *Am J Surg Pathol* 28:748–755.
- Viardot A, Möller P, Högel J, Werner K, Mechttersheimer G, Ho AD, et al. (2002): Clinicopathologic correlations of genomic gains and losses in follicular lymphoma. *J Clin Oncol* 20:4523–4530.
- Vieites B, Fraga M, Lopez-Presas E, Pintos E, Garcia-Rivero A, Forteza J (2005): Detection of t(14;18) translocation in a case of intravascular large B-cell lymphoma: a germinal centre cell origin in a subset of these lymphomas? *Histopathology* 46:466–468.
- Visco C, Tzankov A, Xu-Monette ZY, Miranda RN, Tai YC, Li Y, et al. (2013): Patients with diffuse large B-cell lymphoma of germinal center origin with BCL2 translocations have poor outcome, irrespective of MYC status: a report from an International DLBCL rituximab-CHOP Consortium Program Study. *Haematologica* 98:255–263.
- Walker BA, Wardell CP, Johnson DC, Kaiser MF, Begum DB, Dahir NB, et al. (2013): Characterization of IGH locus breakpoints in multiple myeloma indicates a subset of translocations appear to occur in pregerminal center B-cells. *Blood* 121:3413–3419.
- Wessendorf S, Barth TF, Viardot A, Mueller A, Kestler HA, Kohlhammer H, et al. (2007): Further delineation of chromosomal consensus regions in primary mediastinal B-cell lymphomas: an analysis of 37 tumor samples using high-resolution genomic profiling (array-CGH). *Leukemia* 21:2463–2469.
- Willis TG, Dyer MJ (2000): The role of immunoglobulin translocations in the pathogenesis of B-cell malignancies. *Blood* 96:808–822.
- Willis TG, Jadayel DM, Du MQ, Peng H, Perry AR, Abdul-Rauf M, et al. (1999): Bcl10 is involved in t(1;14)(p22;q32) of MALT B cell lymphoma and mutated in multiple tumor types. *Cell* 96:35–45.
- Wilson KS, McKenna RW, Kroft SH, Dawson DB, Ansari Q, Schneider NR (2002): Primary effusion lymphomas exhibit complex and recurrent cytogenetic abnormalities. *Br J Haematol* 116:113–121.
- Wlodarska I, Matthews C, Veyt E, Pospisilova H, Catherwood MA, Poulsen TS, et al. (2007): Telomeric IGH losses detectable by fluorescence in situ hybridization in chronic lymphocytic leukemia reflect somatic VH recombination events. *J Mol Diagn* 9:47–54.
- Woessmann W (2013): How to treat children and adolescents with relapsed non-Hodgkin lymphoma? *Hematol Oncol Suppl* 1:64–68.
- Wolff DJ, Bagg A, Cooley LD, Dewald GW, Hirsch BA, Jacky PB, et al. (2007): Guidance for fluorescence in situ hybridization testing in hematologic disorders. *J Mol Diagn* 9:134–143.
- Xu WS, Liang RH, Srivastava G (2000): Identification and characterization of BCL6 translocation partner genes in primary gastric high-grade B-cell lymphoma: heat shock protein 89 alpha is a novel fusion partner gene of BCL6. *Genes Chromosomes Cancer* 27:69–75.
- Yan J, Nie K, Mathew S, Tam Y, Cheng S, Knowles DM, et al. (2014): Inactivation of BANK1 in a novel IGH-associated translocation t(4;14)(q24;q32) suggests a tumor suppressor role in B-cell lymphoma. *Blood Cancer J* 4:e215.
- Yates LR, Campbell PJ (2012): Evolution of the cancer genome. *Nat Rev Genet* 13:795–806.
- Ye BH, Lista F, Lo Coco F, Knowles DM, Offit K, Chaganti RS, et al. (1993a): Alterations of a zinc finger-encoding gene, BCL-6, in diffuse large-cell lymphoma. *Science* 262:747–750.
- Ye BH, Rao PH, Chaganti RS, Dalla-Favera R (1993b): Cloning of bcl-6, the locus involved in chromosome translocations affecting band 3q27 in B-cell lymphoma. *Cancer Res* 53:2732–2735.
- Yoshida S, Kaneita Y, Aoki Y, Seto M, Mori S, Moriyama M (1999): Identification of heterologous partner genes fused to the BCL6 gene in diffuse large B-cell lymphomas: 5'-RACE and LA-PCR analyses of biopsy samples. *Oncogene* 18:7994–7999.
- Yunis JJ, Oken MM, Theologides A, Howe RB, Kaplan ME (1984): Recurrent chromosomal defects are found in most patients with non-Hodgkin's-lymphoma. *Cancer Genet Cytogenet* 13:17–28.
- Zani VJ, Asou N, Jadayel D, Heward JM, Shipley J, Nacheva E, et al. (1996): Molecular



cloning of complex chromosomal translocation t(8;14;12)(q24.1;q32.3;q24.1) in a Burkitt lymphoma cell line defines a new gene (BCL7A) with homology to caldesmon. *Blood* 87:3124–3134.

Zech L, Haglund U, Nilsson K, Klein G (1976): Characteristic chromosomal abnormalities in biopsies and lymphoid-cell lines from patients with Burkitt and non-Burkitt lymphomas. *Int J Cancer* 17:47–56.

Zhang S, Zech L, Klein G (1982): High-resolution analysis of chromosome markers in Burkitt lymphoma cell lines. *Int J Cancer* 29:153–157.

Zhang Q, Siebert R, Yan M, Hinzmann B, Cui X, Xue L, et al. (1999): Inactivating mutations and overexpression of BCL10, a caspase recruitment domain-containing gene, in MALT lymphoma with t(1;14)(p22;q32). *Nat Genet* 22:63–68.

Zhang J, Grubor V, Love CL, Banerjee A, Richards KL, Mieczkowski PA, et al. (2013): Genetic heterogeneity of diffuse large B-cell lymphoma. *Proc Natl Acad Sci U S A* 110:1398–1403.

Zinkel S, Gross A, Yang E (2006): BCL2 family in DNA damage and cell cycle control. *Cell Death Differ* 13:1351–1359